

On countable and uncountable sets

Vassos Hadzilacos

A set A is **countable** or **enumerable** if it is finite or there is a bijection $f : \mathbb{N} \rightarrow A$ (i.e., A can be placed in a one-to-one correspondence with the natural numbers). A set is **uncountable** if it is not countable.

From the definition of countable set it follows immediately that A is countable if and only if there is a finite or infinite sequence $S = a_0, a_1, a_2, \dots$ such that every element of A appears once and only once in S . We call such a sequence an **enumeration** of the set A .

In lecture we saw examples of countable and uncountable sets. In particular, we saw two proof techniques, both due to Cantor: **Dovetailing** (merging countably many infinite sequences into a single infinite sequence), which can be used to prove that a set is countable; and **diagonalization**, which can be used to prove that a set is not countable.

Following is a list of the results of this nature that we have proved using these techniques.

Theorem 1.1: \mathbb{Z} is countable.

Theorem 1.2: $\mathbb{N} \times \mathbb{N}$ is countable.

Theorem 1.3: For every positive integer k , $\mathbb{N}^k = \underbrace{\mathbb{N} \times \mathbb{N} \times \dots \times \mathbb{N}}_{k \text{ times}}$ is countable.

Theorem 1.4: $\mathbb{N}^* = \bigcup_{k \in \mathbb{N}} \mathbb{N}^k$ (the set of **finite** sequences of natural numbers) is countable.

Theorem 1.5: The set of infinite sequences over $\{0, 1\}$ is uncountable.

Theorem 1.6: The set of functions from \mathbb{N} to \mathbb{N} is uncountable.

Theorem 1.7: The set of (finite) strings over a (finite) alphabet is countable.

We can leverage the fact that we know that certain sets are countable (or uncountable) to show that other sets have this property.

For example, consider the set of “step functions” from \mathbb{N} to \mathbb{N} . A function $f : \mathbb{N} \rightarrow \mathbb{N}$ is called a **step function** if there is some $n_f \in \mathbb{N}$ and $a, b \in \mathbb{N}$ such that for all n , $0 \leq n \leq n_f$, $f(n) = a$ and for all $n > n_f$, $f(n) = b$. Is the set of step functions from \mathbb{N} to \mathbb{N} countable or uncountable?

Notice that a step function $f : \mathbb{N} \rightarrow \mathbb{N}$ can be uniquely and fully described by a triple of natural numbers (n_f, a, b) . Sometimes we say that this triple **encodes** the function f : Given a step function there is a unique triple that corresponds to it; and conversely, given a triple, there is a unique step function defined by it. Since triples of natural numbers can be encoded by (correspond one-to-one to) natural numbers (see Theorem 1.3), we conclude that

Theorem 1.8: The set of step functions from \mathbb{N} to \mathbb{N} is countable.

The preceding argument exemplifies a useful way to prove that a set is countable: Show that every element of the set can be encoded by (i.e., placed into a one-to-one correspondence with) the elements of a set we have already proved is countable. Now, we can use Theorem 1.8 to show:

Theorem 1.9: The set of non-step functions from \mathbb{N} to \mathbb{N} is uncountable.

PROOF. Suppose, for contradiction, that the set of non-step $\mathbb{N} \rightarrow \mathbb{N}$ functions is countable, and let f_0, f_1, f_2, \dots be an enumeration of this set. By Theorem 1.8, the set of $\mathbb{N} \rightarrow \mathbb{N}$ step functions is countable,

so let g_0, g_1, g_2, \dots be an enumeration of this set. Then we can enumerate the set of all $\mathbb{N} \rightarrow \mathbb{N}$ functions by dovetailing as $f_0, g_0, f_1, g_1, f_2, g_2, \dots$, contradicting Theorem 1.6. \square

As an exercise, try to prove Theorem 1.9 using diagonalization.

Following are some useful facts about countable sets:

Theorem 1.10:

- (a) If $f : \mathbb{N} \rightarrow A$ is an onto (surjective) function, then A is countable.
- (b) If $A \subseteq B$ and B is countable, then A is countable.
- (c) If A and B are countable, then $A \cup B$ is countable.
- (d) If for each $i \in \mathbb{N}$ the set A_i is countable then the set $\bigcup_{i \in \mathbb{N}} A_i$ is countable. (That is, the union of countably many countable sets is countable!).

PROOF. Part (a) is left as an exercise. (See Question 1(a) in Assignment 1!) It implies that if a_0, a_1, a_2, \dots is a sequence of elements of A such that every element of A appears at least once (i.e., this sequence is a listing of all elements of A but possibly with duplicates) then A is countable. This is useful because it is sometimes simpler to describe a listing of a set that may contain duplicates. (Note that the sequence a_0, a_1, a_2, \dots is not an enumeration of A , which is a sequence in which every element of A must appear *exactly* once.)

Part (b) follows from (a): Intuitively, given an enumeration of B we can write an enumeration of A with possible duplicates by replacing elements in the enumeration of B that are not in A by some element of A . (If A is empty, then it is by definition countable since the empty set is finite.)

For part (c), let a_0, a_1, a_2, \dots be an enumeration of A and b_0, b_1, b_2, \dots be an enumeration of B . By dovetailing these two sequences

$$a_0, b_0, a_1, b_1, a_2, b_2, \dots$$

we get an enumeration (with possible repetition) of $A \cup B$. By part (a), $A \cup B$ is countable.

For part (d) we can assume, without loss of generality, that every A_i is non-empty. (If A is countable then $A \cup \emptyset = A$ is also countable.) Let $a_{i0}, a_{i1}, a_{i2}, \dots$ be an enumeration of A_i ; if A_i is finite, we pad the enumeration by repeating some element of A_i an infinite number of times (we can do this because we assume that A_i is non-empty). So, every element of $\bigcup_{i \in \mathbb{N}} A_i$ appears (at least once) in the following table:

a_{00}	a_{01}	a_{02}	a_{03}	\dots
a_{10}	a_{11}	a_{12}	a_{13}	\dots
a_{20}	a_{21}	a_{22}	a_{23}	\dots
a_{30}	a_{31}	a_{32}	a_{33}	\dots
\vdots	\vdots	\vdots	\vdots	\ddots

By dovetailing we can enumerate (possibly with repetition) all elements of all A_i 's. Thus, by part (a), $\bigcup_{i \in \mathbb{N}} A_i$ is countable. \square