

## RIGOROUS HIGH-DIMENSIONAL SHADOWING USING CONTAINMENT: THE GENERAL CASE

CARMEN YOUNG, WAYNE B. HAYES, KENNETH R. JACKSON

Corresponding Author:  
Wayne B Hayes  
Dept. of Computer Science  
University of California, Irvine  
Irvine, California 92697-3425

**Abstract.** A *shadow* is an exact solution to an iterated map that remains close to an approximate solution for a long time. An elegant geometric method for proving the existence of shadows is called *containment*, and it has been proven previously in two and three dimensions, and in some special cases in higher dimensions. This paper presents the general proof using tools from differential and algebraic topology and singular homology.

### 1. Introduction.

1.1. **Background.** An *orbit* of a continuous map  $\varphi$  is a finite or infinite sequence of points generated using

$$\mathbf{x}_{i+1} = \varphi(\mathbf{x}_i). \quad (1)$$

Often one point,  $\mathbf{x}_0$ , is given, called the *initial condition*. Consider an approximation  $\hat{\varphi}$  to  $\varphi$  with the property that

$$\|\hat{\varphi}(\mathbf{x}) - \varphi(\mathbf{x})\| < \delta. \quad (2)$$

An orbit of  $\hat{\varphi}$  generated using

$$\mathbf{y}_{i+1} = \hat{\varphi}(\mathbf{y}_i) \quad (3)$$

is called a  $\delta$ -*pseudo-orbit* of  $\varphi$  and, from (2), has the property

$$\|\mathbf{y}_{i+1} - \varphi(\mathbf{y}_i)\| < \delta \text{ for all } i.$$

Pseudo-orbits are of interest to those studying computer-generated orbits because finite-precision arithmetic is used to compute them, with the consequence that an exact orbit and a pseudo-orbit starting at the same point can diverge exponentially away from each other. See for example Grebogi, Hammel, Yorke, and Sauer (1990), hereafter GHYS. Given a pseudo-orbit (3), the exact orbit (1) is a *shadow* of (3) if

$$\|\mathbf{y}_i - \mathbf{x}_i\| < \varepsilon \text{ for all } i.$$

Shadowing was first discussed by Anosov (1967) and Bowen (1975), in relation to *hyperbolic* systems, in which space along an orbit can be uniformly separated into *expanding* and *contracting* subspaces. Let  $S$  and  $\varphi$  be the invariant set and the map of a hyperbolic system, respectively. In such systems, Anosov (1967) proved that  $\forall \varepsilon > 0, \exists \delta > 0$  such that every infinite-length  $\delta$ -pseudo orbit remaining in  $S$  is  $\varepsilon$ -shadowed by an exact trajectory in  $S$ . Bowen (1975) proved that the same result

---

1991 *Mathematics Subject Classification*. Primary: Secondary:

*Key words and phrases*. Nonlinear dynamical systems, shadowing, reliable simulation.

This research was supported by NSERC Canada.




FIGURE 1. Containment in 3D with 2 expanding directions and 1 contracting.

holds if the map is required to be hyperbolic only along trajectories in the vicinity of the pseudo-orbit. Palmer (1988) proved a similar theorem along the way towards using the theory of exponential dichotomies to prove Smale's Theorem (Smale 1965, 1967).

Most systems of general interest, however, are not hyperbolic. The first studies of shadows for non-hyperbolic systems appear to be Beyn (1987) and Hammel, Yorke, and Grebogi (1987). Hammel, Yorke, and Grebogi (1988) and GHYS provide the first proof of the existence of a shadow for a two-dimensional non-hyperbolic system over a non-trivial length of time, using a method called *containment*. Here, by way of introduction, we outline a three-dimensional case that is proved in Hayes and Jackson (2003).

Let  $\varphi$  be a map which is not hyperbolic, but which displays *pseudo-hyperbolicity* (Hayes 2001) for a finite but non-trivial number of iterations. Let  $\{y_i\}_{i=a}^b \subset \mathbb{R}^3$  be a three-dimensional  $\delta$ -pseudo-orbit of  $\varphi$  for integers  $a$  and  $b$ . In this case the pseudo-orbit has 1 contracting direction and two expanding directions (Figure 1), and pseudo-hyperbolicity means that as  $i$  increases, orbits separated from each other by a small distance in the expanding subspace diverge on average (but not necessarily uniformly) away from each other, while orbits separated by a small distance in the contracting subspace approach each other, on average. The three-dimensional containment process consists of building a parallelogram  $M_i$  around each point  $y_i$  of the pseudo-orbit such that the first pair of expanding faces  $F_i^{\pm 1}$  are separated along one expanding direction (the  $x$  direction in Figure 1), the second pair of expanding faces  $F_i^{\pm 2}$  are separated along the other expanding direction (the  $y$  direction in Figure 1), and the one pair of contracting faces  $F_i^{\pm 3}$  are separated from each other along the contracting direction (the  $z$  direction in Figure 1). In order to prove the existence of a shadow, we require that  $\varphi(M_i)$  maps over  $M_{i+1}$  so that  $\varphi$  flattens  $M_i$  into a thin slice, cutting  $M_{i+1}$  into 3 pieces, the middle piece of which contains a contiguous section of  $\varphi(M_i)$  (as well as possibly some isolated pieces of  $\varphi(M_i)$ ). Now, assume  $\gamma_i$  is a surface in  $M_i$  whose boundary connects and “wraps around” all of the expanding sides of  $M_i$ . Then there is a contiguous patch of  $\varphi(\gamma_i) \cap M_{i+1} \equiv \gamma_{i+1}$  lying wholly in  $M_{i+1}$  whose boundary  $\partial\gamma_{i+1}$  connects and “wraps around” the expanding sides of  $M_{i+1}$ . If this property continues for each step then, by induction, there is a  $\gamma_N \neq \emptyset$  lying wholly within  $M_N$  whose boundary  $\partial\gamma_N$  connects and wraps around the expanding sides of  $M_N$ . Then any point  $x_N \in \gamma_N$

can be traced backwards to a point  $\mathbf{x}_i \in \gamma_i \subset M_i$  for  $i = 0, 1, \dots, N-1$ , and the  $\mathbf{x}_i$  trajectory is an exact orbit lying close the pseudo-orbit — *i.e.*, a shadow.

This case, along with all other one-, two-, and three-dimensional cases, as well as some special cases in higher dimension, were proved in Hayes and Jackson (2003). The purpose of this paper is to present the general  $n$ -dimensional proof in which  $k$  directions are expanding, while  $n - k$  directions are contracting.

**1.2. Overview.** The machinery that we use requires that the intersections of the manifolds  $\varphi(\gamma_i) \cap M_{i+1}$  are transversal. Theorem 6 (Sard's Theorem) demonstrates that there exists  $\gamma_0$  such that for every  $i > 0$ ,  $\varphi(\gamma_i)$  is transversal to  $M_{i+1}$ . In Section 2 we present the main result. Section 3 presents the background for Sard's Theorem, while Section 4 provides a brief background to singular homology and cohomology, and finally, the proof of our main result.

**2. Main Result.** Let  $\varphi : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be a diffeomorphism with  $k$  expanding directions and  $(n - k)$  contracting directions. For each  $i = 0, \dots, N$ , let  $M_i$  be an  $n$ -cube in  $\mathbb{R}^n$ . We assume that the faces of  $M_i$  are labeled so that the first  $2k$  faces  $F_i^{\pm j}$ ,  $j = 1, \dots, k$ , lie transverse to the expanding directions of  $\varphi$ , and the remaining  $2(n - k)$  faces  $F_i^{\pm j}$ ,  $j = k + 1, \dots, n$ , lie transverse to the contracting directions of  $\varphi$ . We denote the union of a set of faces by listing multiple integers in the superscript. Thus the *expanding faces* of  $M_i$  are collectively denoted  $\partial_X M_i \equiv F_i^{\pm 1, \dots, \pm k}$  and the *contracting faces* of  $M_i$  are denoted  $\partial_C M_i \equiv F_i^{\pm(k+1), \dots, \pm n}$ .

Let  $\text{Int}(A)$  denote the interior of the set  $A$ . Refer to Figure 2. We say that  $M_i$  and  $M_{i+1}$  satisfy the  $(n, k)$ -*Inductive Containment Property* (abbreviated  $(n, k)$ -ICP), for  $\varphi$  if

- (ICP1)  $\varphi(\partial_X M_i) \cap M_{i+1} = \emptyset$  and, for all  $j \in \{1, \dots, k\}$ ,  $\varphi(F_i^{-j})$  and  $\varphi(F_i^{+j})$  lie on opposite sides of the infinite slab between the two hyperplanes containing  $F_{i+1}^{-j}$  and  $F_{i+1}^{+j}$ , respectively.
- (ICP2) There is a parallelepiped  $Q_{i+1} \subset \mathbb{R}^n$  with faces  $G_{i+1}^j$  parallel to the faces  $F_{i+1}^j$  of  $M_{i+1}$  for  $j = \pm 1, \dots, \pm n$  such that
  - i)  $\varphi(M_i) \subset \text{Int}(Q_{i+1})$ ,
  - ii)  $Q_{i+1} \cap \partial_C M_{i+1} = \emptyset$  and, for all  $j \in \{k + 1, \dots, n\}$ ,  $F_{i+1}^{-j}$  and  $F_{i+1}^{+j}$  lie on opposite sides of the infinite slab between the two hyperplanes containing  $G_{i+1}^{-j}$  and  $G_{i+1}^{+j}$ .

The conditions of the Inductive Containment Property can be rigorously verified computationally (Hayes 2001).

**Theorem 1** ( $(n, k)$ -Inductive Containment Theorem). *Suppose that  $M_i$  and  $M_{i+1}$  satisfy  $(n, k)$ -ICP for  $\varphi$  for all  $i = 0, \dots, N - 1$ . Then there exists a sequence of non-empty  $k$ -manifolds  $\gamma_i \subset M_i$ ,  $i = 0, \dots, N$ , such that*

$$\text{Int}(\gamma_i) \subset \text{Int}(M_i), \quad \partial\gamma_i = \gamma_i \cap \partial_X M_i, \quad \text{and} \quad \gamma_{i+1} \subset \varphi(\gamma_i)$$

for all  $i$ .

We will prove this theorem in stages. For a cleaner exposition of the proof, we will translate all objects to a *standardized* frame in the vicinity of the origin, as follows. Let  $\square_n$  denote the standard unit cube in  $\mathbb{R}^n$ ,

$$\square_n = \{(x_1, \dots, x_n) : |x_j| \leq 1 \text{ for } j = 1, \dots, n\},$$




FIGURE 2. Schematic diagram of the Inductive Containment Property in 2 dimensions. The tall parallelogram is  $Q_{i+1} \supset \varphi(M_i)$ , the wider one is  $M_{i+1}$ . The vertical direction is expanding. The horizontal direction is contracting.

and denote its faces by

$$E^{+j} = \{(x_1, \dots, x_n) : x_j = +1, |x_l| \leq 1 \text{ for } l \neq j\}, \quad j = 1, \dots, n,$$

$$E^{-j} = \{(x_1, \dots, x_n) : x_j = -1, |x_l| \leq 1 \text{ for } l \neq j\}, \quad j = 1, \dots, n.$$

We introduce the *Standardized  $(n, k)$ -Inductive Containment Property* by transforming both  $M_i$  and  $M_{i+1}$  to  $\square_n$ , as follows. For each  $i = 0, \dots, N$  there is an orientation-preserving diffeomorphism (*i.e.*, change of coordinates)  $\psi_i : \mathbb{R}^n \rightarrow \mathbb{R}^n$  that maps  $M_i$  to  $\square_n$ , and maps  $F_i^{\pm j}$  to  $E^{\pm j}$  for  $j = 1, \dots, n$ . Let  $\mathcal{M}_i = \square_n$  for all  $i$ . Let  $\phi_i = \psi_{i+1} \circ \varphi \circ \psi_i^{-1}$ . If  $M_i$  and  $M_{i+1}$  satisfy  $(n, k)$ -ICP for  $\varphi$ , then by construction  $\mathcal{M}_i$  and  $\mathcal{M}_{i+1}$  satisfy  $(n, k)$ -ICP for  $\phi_i$ , and we say that the Standardized  $(n, k)$ -ICP holds for  $\phi_i$ . Note that it is easy to choose  $\psi_i$  so that the Standardized  $(n, k)$ -ICP holds for  $\phi$ , such that there exists positive  $\varepsilon < 1$  such that (ICP2) holds with

$$Q = \{(x_1, \dots, x_n) : |x_i| \leq 1/\varepsilon \text{ for } i = 1, \dots, k \text{ and } |x_i| \leq 1 - \varepsilon \text{ for } i = k+1, \dots, n\}.$$

For our purposes, the term *manifold* will refer to a smooth manifold with boundary and corners.

**Definition 1.** Suppose that  $\Gamma \subset \square_n$  is a  $k$ -manifold with  $\partial\Gamma \subset \partial_X \square_n$ . We say that  $\partial\Gamma$  **wraps around**  $\partial_X \square_n$  if the homology class of  $[\partial\Gamma]$  in  $H_{k-1}(\partial_X \square_n)$  is **not** zero.

**Lemma 1.** Let  $\phi : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be a diffeomorphism and assume that the standardized  $(n, k)$ -ICP holds for  $\phi$ . Let  $\Gamma \subset \square_n$  be a non-empty  $k$ -manifold with boundary  $\partial\Gamma \subset \partial_X \square_n$ , and suppose further that  $\partial\Gamma$  wraps around  $\partial_X \square_n$ . Finally, suppose that  $\phi(\Gamma)$  is transverse to  $\partial_X \square_n$ , and let  $\Gamma' \equiv \phi(\Gamma) \cap \square_n$ . Then the following hold:

- i)  $\Gamma'$  is a non-empty  $k$ -manifold with boundary  $\partial\Gamma' = \Gamma' \cap \partial \square_n$ .
- ii)  $\partial\Gamma' \subset \partial_X \square_n$ .
- iii)  $\partial\Gamma'$  wraps around  $\partial_X \square_n$ .

We will prove this lemma in Section 4.

*Proof of Theorem 1.* We will prove the theorem by induction. Let  $\psi_i : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be a change of co-ordinates that maps  $M_i$  to  $\square_n$ , and maps  $F_i^{\pm j}$  to  $E^{\pm j}$  for  $j = 1, \dots, n$ . Let  $\phi_i = \psi_{i+1} \circ \varphi \circ \psi_i^{-1}$ . Then the Standardized  $(n, k)$ -ICP holds for each  $\phi_i$ . Let  $\square_k$  denote the unit  $k$ -cube in  $\mathbb{R}^n$ ,

$$\square_k = \{(x_1, \dots, x_n) \in \mathbb{R}^n : |x_i| \leq 1 \text{ for } 1 \leq i \leq k, x_{k+1} = \dots = x_n = 0\}.$$

This is a  $k$ -dimensional submanifold of  $\square_n$ , and its boundary

$$\partial \square_k = \{(x_1, \dots, x_n) \in \square_k : x_j = \pm 1 \text{ for some } j \in \{1, \dots, k\}\},$$

is contained in  $\partial_X \square_n$ . By Lemma 3 and Definition 6, the homology class  $[\partial \square_k] \in H_{k-1}(\partial_X \square_n)$  is 1. By Theorem 6 (Sard's Theorem), we can homotope  $\square_k$  relative to its boundary to a  $k$ -manifold  $\Gamma$  with  $\partial \Gamma = \partial \square_k$  such that  $\Gamma$  intersects  $\phi_0^{-1} \phi_1^{-1} \dots \phi_i^{-1}(\partial_X \square_n)$  transversally for each  $i = 0, \dots, N$ . Thus  $[\partial \Gamma] = [\partial \square_k] \neq 0$ , and

$$\phi_i \phi_{i-1} \dots \phi_0(\Gamma) \text{ is transverse to } \partial_X \square_n \text{ for all } i. \quad (4)$$

We start the induction by taking  $\Gamma_0 = \Gamma$ . Then  $\Gamma_0$  and  $\phi_0$  satisfy the hypotheses of Lemma 1. Let  $\Gamma_1 = \phi_0(\Gamma_0) \cap \square_n$ . Then  $\Gamma_1$  is a non-empty  $k$ -manifold with boundary  $\partial \Gamma_1 \subset \partial_X \square_n$ , and  $[\partial \Gamma_1] = [\partial \Gamma_0] \neq 0$ .

At the  $i$ -th step of the induction, we have a non-empty  $k$ -manifold  $\Gamma_i \subset \square_n$  with  $\partial \Gamma_i \subset \partial_X \square_n$  and  $[\partial \Gamma_i] \neq 0$  in  $H_{k-1}(\partial_X \square_n)$ . Moreover  $\Gamma_i \subset \phi_{i-1}(\Gamma_{i-1}) \subset \dots \subset \phi_{i-1} \phi_{i-2} \dots \phi_0(\Gamma_0)$ , which implies that  $\phi_i(\Gamma_i) \subset \phi_i \dots \phi_0(\Gamma_0)$ . From (4), we see that  $\phi_i(\Gamma_i)$  is transverse to  $\partial_X \square_n$ , so we can apply Lemma 1. If we set  $\Gamma_{i+1} = \phi_i(\Gamma_i) \cap \square_n$ , then  $\Gamma_{i+1}$  and  $\phi_{i+1}$  satisfy the hypotheses of Lemma 1, and the induction continues.

By the  $N$ -th step of the induction, we have produced non-empty  $k$ -manifolds  $\Gamma_i$ ,  $i = 0, \dots, N$ , with  $\text{Int}(\Gamma_i) \subset \text{Int}(\square_n)$ ,  $\partial \Gamma_i = \Gamma_i \cap \partial_X \square_n$ ,

$$[\partial \Gamma_i] = [\partial \square_k] \neq 0 \text{ for all } i,$$

and

$$\Gamma_{i+1} \subset \phi_i(\Gamma_i) \text{ for all } i.$$

Let  $\gamma_i = \psi_i^{-1}(\Gamma_i)$  for each  $i = 0, \dots, N$ . Then  $\text{Int}(\gamma_i) \subset \text{Int}(M_i)$ ,  $\partial \gamma_i = \gamma_i \cap \partial_X M_i$ , and

$$\gamma_{i+1} = \psi_{i+1}^{-1}(\Gamma_{i+1}) \subset \psi_{i+1}^{-1}(\phi_i(\Gamma_i)) = \psi_{i+1}^{-1}(\psi_{i+1} \varphi \psi_i^{-1})(\Gamma_i) = \varphi(\gamma_i).$$

This completes the proof of Theorem 1.

**Corollary 1** (Shadowing Containment Theorem). *Let  $\{M_i\}_{i=0}^N$  be a sequence of  $n$ -cubes enclosing a pseudo-trajectory  $\{\mathbf{y}_i\}_{i=0}^N$ , and suppose that  $M_i$  and  $M_{i+1}$  satisfy  $(n, k)$ -ICP for all  $i$ . Let  $\varepsilon$  be the maximum diameter of  $M_i$  over all  $i$ . Then there exists an  $\varepsilon$ -shadow  $\{\mathbf{x}_i\}_{i=0}^N$  of  $\{\mathbf{y}_i\}_{i=0}^N$ , i.e.,  $\|\mathbf{x}_i - \mathbf{y}_i\| \leq \varepsilon$  for all  $i = 0, \dots, N$ .*

*Proof.* By the Inductive Containment Theorem, there is a sequence of non-empty  $k$ -manifolds  $\gamma_i \subset M_i$  such that  $\gamma_i \subset \text{Int}(\varphi(\gamma_{i-1}))$  for all  $i$ . Let  $\mathbf{x}_N$  be any point in  $\gamma_N$ . As  $\varphi$  is a diffeomorphism, there is a unique point  $\mathbf{x}_0 \in \gamma_0$  such that  $\varphi^N(\mathbf{x}_0) = \mathbf{x}_N$ . For each  $i$  we set  $\mathbf{x}_i = \varphi^i(\mathbf{x}_0)$ . Then  $\mathbf{x}_{i+1} = \varphi(\mathbf{x}_i) \in M_{i+1}$  for all  $i$  and  $\{\mathbf{x}_i\}_{i=0}^N$  is an  $\varepsilon$ -shadow of  $\{\mathbf{y}_i\}_{i=0}^N$ .  $\square$

**3. Sard's Theorem and Transversality.** For the purpose of proving the Inductive Containment Theorem in arbitrary dimensions, we would like to determine under what conditions two objects in  $\mathbb{R}^n$  intersect in a "nice" way. It turns out that when the objects in question are smooth manifolds, then a good answer is provided by *transversality theory*. In this section we review some of the basic concepts from transversality theory that we need. Guillemin and Pollack (1974) supply a more detailed introduction, together with some applications to geometry.

**3.1. Manifolds.** Let  $X$  be a smooth manifold of dimension  $k$ . That is, every point  $x \in X$  has an open neighborhood  $V$  which is homeomorphic to an open set  $U \subset \mathbb{R}^k$ :

$$\theta : U \xrightarrow{\cong} V \subset X. \quad (5)$$

The triplet  $(\theta, U, V)$  is sometimes referred to as a *coordinate chart* near  $x$ . If  $(\theta_1, U_1, V_1)$ ,  $(\theta_2, U_2, V_2)$  are two coordinate charts (near points  $x_1$  and  $x_2$ , say) that *overlap* in the sense that  $V_1 \cap V_2 \neq \emptyset$ , then we further require that

$$\begin{aligned} \theta_1^{-1} \circ \theta_2 : \theta_2^{-1}(V_1 \cap V_2) &\rightarrow \theta_1^{-1}(V_1 \cap V_2) \\ \text{and } \theta_2^{-1} \circ \theta_1 : \theta_1^{-1}(V_1 \cap V_2) &\rightarrow \theta_2^{-1}(V_1 \cap V_2) \end{aligned}$$

be smooth, as maps between open sets in  $\mathbb{R}^k$ .

**3.2. Tangent space.** In the cases of interest to us, the manifold  $X$  sits in some higher-dimensional Euclidean space  $\mathbb{R}^n$ , so the coordinate map (5) is simply a smooth map from  $U \subset \mathbb{R}^k$  to  $\mathbb{R}^n$ , with image equal to  $V$ . Thus the derivative of  $\theta$  at a point  $u \in U$  makes sense as a linear map from  $\mathbb{R}^k$  to  $\mathbb{R}^n$ , *i.e.*, it is a real  $n \times k$  matrix. We denote this derivative by  $d\theta_u$ .

Suppose  $(\theta, U, V)$  is a coordinate chart near  $x \in X$ , and  $\theta(u) = x$ . We define the *tangent space to  $X$  at  $x$*  to be the image of  $d\theta_u : \mathbb{R}^k \rightarrow \mathbb{R}^n$ . (Equivalently, the tangent space is the linear span of the column vectors of  $d\theta_u$ .) This is a vector subspace of  $\mathbb{R}^n$  which we denote by  $T_x X$ . Geometrically,

$$x + T_x X = \{x + v \mid v \in T_x X\}$$

consists of all vectors starting at  $x$  that are tangent there to  $X$ . In other words, it is the best approximation of  $X$  near  $x$  by a linear subspace. One can easily check that if  $(\theta', U', V')$  is another coordinate chart near  $x$  with  $\theta'(u') = x$ , then  $\text{Image}(d\theta_u) = \text{Image}(d\theta'_{u'})$  – that is,  $T_x X$  is well-defined and independent of our choice of coordinate chart.

**3.3. Smooth maps and differentials.** If  $X^k$  and  $Y^l$  are smooth manifolds of respective dimensions  $k$  and  $l$ , then a continuous map  $f : X \rightarrow Y$  is called *smooth* if, for every coordinate chart  $(\theta, U, V)$  for  $X$  and  $(\tilde{\theta}, \tilde{U}, \tilde{V})$  for  $Y$ , the restriction

$$\tilde{\theta}^{-1} \circ f \circ \theta|_{(f \circ \theta)^{-1}(\tilde{U})}$$

is smooth when considered as a map defined in an open subset of  $\mathbb{R}^k$  with values in  $\mathbb{R}^l$ . If  $X \subset \mathbb{R}^m$  and  $Y \subset \mathbb{R}^n$  for some  $m > k$ ,  $n > l$ , it is equivalent to say that  $f : X \rightarrow Y$  is smooth if around any point  $x \in X$  there is an open ball  $B_x$  and a smooth map  $F : B_x \rightarrow \mathbb{R}^n$  such that the restriction of  $F$  to  $X \cap B_x$  equals  $f$ :

$$F|_{X \cap B_x} = f.$$

If  $f$  is smooth and  $f(x) = y$ , then the *derivative* of  $f$  at  $x$  is a linear map

$$df_x : T_x X \rightarrow T_y Y.$$

To define the derivative, suppose that  $(\theta, U, V)$  is a coordinate chart near  $x$  with  $\theta(u) = x$ , and  $(\tilde{\theta}, \tilde{U}, \tilde{V})$  is a coordinate chart near  $y$  with  $\tilde{\theta}(\tilde{u}) = y$ . Set  $h = \tilde{\theta}^{-1} \circ f \circ \theta$  on  $(f \circ \theta)^{-1}(\tilde{V})$  so that the diagram

$$\begin{array}{ccc} V & \xrightarrow{f} & \tilde{V} \\ \theta \uparrow & & \uparrow \tilde{\theta} \\ U & \xrightarrow{h} & \tilde{U} \end{array}$$

commutes. Note in particular that  $h(u) = \tilde{u}$ . We then set

$$df_x(\nu) = d\tilde{\theta}_{\tilde{u}} \circ dh_u \circ (d\theta_u)^{-1}(\nu), \text{ for } \nu \in \text{Image}(d\theta_u).$$

(If we think of the derivatives of  $\tilde{\theta}, h$  and  $\theta$  as matrices, then the latter composition is a product of matrices.) This clearly maps  $T_x X = \text{Image}(d\theta_u)$  into  $T_y Y = \text{Image}(d\tilde{\theta}_{\tilde{u}})$ . If  $X \subset \mathbb{R}^m$ ,  $Y \subset \mathbb{R}^n$  and  $f$  near  $x$  equals the restriction of  $F : B_x \rightarrow \mathbb{R}^n$ , then it is not hard to show that  $df_x$  equals the restriction of  $dF_x$  to  $T_x X \subset \mathbb{R}^m$ .

If  $x \in X$  and  $df_x = 0$ , that is,  $df_x(\nu) = 0$  for all  $\nu \in T_x X$ , then  $x$  is called a *critical point* of  $f$ . If  $y \in Y$  and there is some  $x \in f^{-1}(y)$  such that  $df_x = 0$ , then  $y$  is called a *critical value* of  $f$ . If no such  $x$  exists for  $y$ , then  $y$  is called a *regular value* of  $f$ .

**3.4. Transversality.** The smooth map  $f : X \rightarrow Y$  is *transversal* to the submanifold  $Z \subset Y$  if

$$\text{Image}(df_x) + T_{f(x)}(Z) = T_{f(x)}Y$$

for every point  $x$  in the preimage of  $Z$ .

**Theorem 2** (Preimage Theorem). *Let  $f : X \rightarrow Y$  be a smooth map of manifolds. If  $f$  is transversal to a submanifold  $Z \subset Y$ , then  $f^{-1}(Z)$  is a submanifold of  $X$  and the codimension of  $f^{-1}(Z)$  in  $X$  equals the codimension of  $Z$  in  $Y$ .*

*Note:* The codimension of  $Z$  in  $Y$  is  $\text{codim } Z = \dim Y - \dim Z$ .

**Definition 2.** Suppose  $X \subset Y$ . The **inclusion map**  $i : X \rightarrow Y$  is defined by  $i(x) = x$  for all  $x \in X$ .

When  $X$  and  $Z$  are both submanifolds of the same manifold  $Y$ , then a point  $x \in X$  lies in the intersection  $X \cap Z$  if and only if  $x$  lies in the preimage of  $Z$  under the inclusion map  $i : X \hookrightarrow Y$ . We say that  $X$  and  $Z$  *intersect transversally* in  $Y$  if

$$T_x X + T_x Z = T_x Y$$

for every  $x \in X \cap Z$ . Since the derivative  $di_x : T_x X \rightarrow T_x Y$  is simply the inclusion of  $T_x X$  into  $T_x Y$ , the next result is a direct consequence of the Preimage Theorem.

**Theorem 3.** *If the submanifolds  $X$  and  $Z$  intersect transversally in  $Y$ , then their intersection  $X \cap Z$  is again a submanifold and*

$$\text{codim}(X \cap Z) = \text{codim } X + \text{codim } Z.$$

**Theorem 4** (Transversality Theorem, (Guillemin and Pollack 1974)). *Let  $F : X \times S \rightarrow Y$  be a smooth map of manifolds. Let  $Z \subset Y$  be a smooth submanifold without boundary. If  $F$  is transverse to  $Z$  then for almost every  $s \in S$ ,  $F_s = F(\cdot, s)$  is transverse to  $Z$ .*

The theorem follows from an application of Sard's theorem, which we now state. (See (Guillemin and Pollack 1974, Chap 2,§1).)

**Theorem 5** (Sard's Theorem). *For any smooth map of a manifold  $X$  (with boundary) into a boundaryless manifold  $Y$ , almost every point of  $Y$  is a regular value of  $f : X \rightarrow Y$  (and of  $\partial f = f|_{\partial X} : \partial X \rightarrow Y$ ).*

The idea behind the proof of Theorem 4 is this. By the Preimage theorem,  $W = F^{-1}(Z)$  is a submanifold of  $X \times S$ . Let  $\pi : X \times S \rightarrow S$  be the natural projection map, and consider its restriction to  $W \subset X \times S$ . By Sard's theorem, almost every value of  $s \in S$  is a regular value of  $\pi : W \rightarrow S$ . Using the fact that  $F$  is transversal to  $Z$ , one can show that the regular values of  $\pi|_W$  correspond to the values of  $s$  for which  $F_s$  is transversal to  $Z$ . For details of the proof, we refer the reader to (Guillemin and Pollack 1974, Chap 2, §3).

Combining the Transversality theorem with the Preimage theorem, one can prove that any smooth map  $X \rightarrow Y$  can be deformed by an arbitrarily small amount to a map that is transversal to a submanifold  $Z \subset Y$ . As a special case of this, we have the following.

*Notation:* Given an open set  $V$ , we write  $U \subset\subset V$  to signify that there is a compact set  $K$  with  $U \subset K \subset V$ .

**Theorem 6.** *Let  $\square_k$  denote the  $k$ -cube*

$$\square_k = \{(x_1, \dots, x_n) \in \mathbb{R}^n : |x_i| \leq 1 \text{ for } 1 \leq i \leq k, x_{k+1} = \dots = x_n = 0\},$$

*with boundary*

$$\partial \square_k = \{(x_1, \dots, x_n) \in \square_k : x_i = \pm 1 \text{ for some } i \in \{1, \dots, k\}\}.$$

*Let  $Z_1, \dots, Z_L$  be smooth submanifolds of  $\mathbb{R}^n$ , and suppose each has the property that  $Z_l \cap \square_k \subset\subset \text{int } \square_k$ . Then we can homotope  $\square_k$  relative to its boundary to a  $k$ -manifold  $\Gamma$  with  $\partial \Gamma = \partial \square_k$ , so that  $\Gamma$  intersects  $Z_l$  transversally for all  $l = 1, \dots, L$ .*

*Proof.* There exists a compact set  $K$  such that  $Z_l \cap \square_k \subset K \subset \text{int } \square_k$  for all  $l$ . Take  $\varepsilon : \square_k \rightarrow \mathbb{R}$  to be a smooth, compactly supported bump function with  $\text{spt}(\varepsilon) \subset \text{int } \square_k$ . We can assume that  $K \subset \{x : \varepsilon(x) \neq 0\}$ .

Let  $S$  denote the open unit ball in  $\mathbb{R}^n$  and let  $F : \square_k \times S \rightarrow \mathbb{R}^n$  be the smooth map  $F(x, s) = x + \varepsilon(x) \cdot s$ . For any fixed point  $x$  where  $\varepsilon(x) \neq 0$ , the map  $s \mapsto F(x, s)$  is a rescaling followed by translation of the ball  $S$ , hence is a submersion. If  $Z \subset \mathbb{R}^n$  is a submanifold and  $Z \cap \square_k \subset \{x : \varepsilon(x) \neq 0\}$ , then it follows that  $F$  is transversal to  $Z$ . So by the transversality theorem of Guillemin-Pollack, the map  $x \mapsto F(x, s)$  is transverse to  $Z$  for almost every  $s \in S$ .

By our hypotheses,  $Z_l \cap \square_k \subset \{x : \varepsilon(x) \neq 0\}$  for every  $l$ . Thus for each  $l$ , there is a subset  $\Omega_l \subset S$  of measure zero such that  $F_s = F(\cdot, s)$  is transverse to  $Z_l$  for any  $s \in S \setminus \Omega_l$ . The union  $\Omega = \Omega_1 \cup \dots \cup \Omega_L$  again has measure zero, and for any  $s \in S \setminus \Omega$ ,  $F_s$  is transverse to all the  $Z_l$ . Now the set  $S \cap \square_k$  also has measure zero and for any  $s \in S \setminus \square_k$ ,

$$\Gamma_s = \{x + \varepsilon(x) \cdot s : x \in \square_k\}$$

is a smooth submanifold of  $\mathbb{R}^n$  with boundary (and corners). Moreover, if  $s \in S \setminus (\Omega \cup \square_k)$ , then saying that  $F_s$  is transverse to  $Z_l$  is equivalent to saying that  $\Gamma_s$  and  $Z_l$  intersect transversally.

Let us fix one such  $s$  and take  $\Gamma = \Gamma_s$  to be our desired  $k$ -manifold. The homotopy from  $\square_k$  to  $\Gamma$  is given by

$$h_t(x) = x + t\varepsilon(x) \cdot s, \quad t \in [0, 1].$$

Clearly  $h_0 : \square_k \rightarrow \square_k$  is the identity map and  $h_1$  maps  $\square_k$  homeomorphically onto  $\Gamma$ . For any  $x \in \partial\square_k$ , we have  $\varepsilon(x) = 0$  because  $\text{spt}(\varepsilon) \subset \text{int}\square_k$ . Therefore  $h_t(x) = x$  for all  $x \in \partial\square_k$  and all  $t \in [0, 1]$ , which is to say that the homotopy fixes the boundary and, in particular,  $\partial\Gamma = \partial\square_k$ .  $\square$

#### 4. Proof of Lemma 1.

**4.1. Review of Singular Homology (with integer coefficients).** For a quick introduction to singular homology (and cohomology), we refer the reader to Appendix A of Milnor and Stasheff (1974). More details can also be found in the graduate text by Greenberg (1981).

The basic objects of singular homology are *equivalence classes* of *singular simplices* in a predetermined topological space. These in turn are modeled on standard simplices in Euclidean space.

**Definition 3.** Let  $K \geq 0$ . The **standard  $K$ -simplex** is the convex set  $\Delta^K \subset \mathbb{R}^{K+1}$ , consisting of all  $(K+1)$ -tuples  $(y_0, \dots, y_K)$  with

$$y_i \geq 0 \quad , \quad y_0 + y_1 + \dots + y_K = 1.$$

Any continuous map  $\sigma$  from  $\Delta^K$  to a topological space  $X$  is called a **singular  $K$ -simplex** in  $X$ .

Let  $K \geq 0$  be an integer and let  $C_K(X)$  be the free  $\mathbb{Z}$ -module obtained by taking one generator  $[\sigma]$  for each singular  $K$ -simplex in  $X$ . We call  $C_K(X)$  the  $K$ -th *singular chain group* of  $X$ . For  $K < 0$ ,  $C_K(X)$  is defined to be zero.

To define the equivalence relation on  $C_K(X)$ , we need to introduce the following *boundary operator*.

**Definition 4.** Let  $\sigma : \Delta^K \rightarrow X$  be a singular  $K$ -simplex in  $X$ . The  $i$ -th *face* of  $\sigma$  is the singular  $(K-1)$ -simplex

$$\sigma \circ \lambda_i : \Delta^{K-1} \rightarrow X,$$

where the linear embedding  $\lambda_i : \Delta^{K-1} \rightarrow \Delta^K$  is defined by

$$\lambda_i(y_0, \dots, y_{i-1}, y_{i+1}, \dots, y_K) = (y_0, \dots, y_{i-1}, 0, y_{i+1}, \dots, y_K).$$

The homomorphism

$$\partial : C_K(X) \rightarrow C_{K-1}(X)$$

given by

$$\partial[\sigma] = [\sigma \circ \lambda_0] - [\sigma \circ \lambda_1] + \dots + (-1)^K [\sigma \circ \lambda_K]$$

is called the **boundary homomorphism**.

It is an exercise in algebra to verify that

$$\partial \circ \partial = 0. \tag{6}$$

Let  $\mathcal{Z}_K(X)$  be the kernel of  $\partial : C_K(X) \rightarrow C_{K-1}(X)$ , and let  $\mathcal{B}_K(X)$  be the image of  $\partial : C_{K+1}(X) \rightarrow C_K(X)$ . By (6),  $\mathcal{B}_K(X) \subset \mathcal{Z}_K(X)$  so the quotient

$$H_K(X) = \mathcal{Z}_K(X) / \mathcal{B}_K(X) \tag{7}$$

makes sense. We call  $H_K(X)$  the  $K$ -th *singular homology group* of  $X$ , and an element of  $H_K(X)$  is called a *homology class*.

Suppose  $f : X \rightarrow Y$  is a continuous map. By composing with  $f$ , we get a map

$$f \circ : \{K\text{-simplices in } X\} \rightarrow \{K\text{-simplices in } Y\}$$

which maps  $\sigma$  to  $f \circ \sigma$ . One can show further that there is an ‘‘induced’’ map

$$f_* : H_K(X) \rightarrow H_K(Y).$$

See Appendix A of Milnor and Stasheff (1974), or Greenberg (1981) for details.

**Definition 5.** We call  $f_* : H_K(X) \rightarrow H_K(Y)$  the **push forward map** of  $f$ .

The following is a basic result in homology theory.

**Proposition 1.** *Let  $f_1, f_2 : X \rightarrow Y$  be continuous maps and suppose that  $f_1$  is homotopic to  $f_2$ . Then the push-forward maps  $(f_1)_* : H_K(X) \rightarrow H_K(Y)$  and  $(f_2)_* : H_K(X) \rightarrow H_K(Y)$  are equal. That is,  $(f_1)_* = (f_2)_*$  as maps from  $H_K(X)$  to  $H_K(Y)$ .*

**Proposition 2.** *Let  $m$  be a positive integer and let  $S^m$  be the  $m$ -dimensional sphere. Then*

$$H_i(S^m) \cong \begin{cases} \mathbb{Z} & \text{if } i = 0 \text{ or } i = m \\ 0 & \text{otherwise.} \end{cases}$$

**Lemma 2.**  $Q \setminus \square_n$  is homotopic to  $S^{k-1}$ .

**Lemma 3.**  $\partial_X \square_n$  is homotopic to  $S^{k-1}$ .

**Lemma 4.** *If  $\phi$  satisfies the Standardized  $(n, k)$ -ICP then  $\phi|_{\partial_X \square_n} : \partial_X \square_n \rightarrow Q \setminus \square_n$  induces an isomorphism in homology, that is,*

$$\phi_* : H_r(\partial_X \square_n) \rightarrow H_r(Q \setminus \square_n)$$

is an isomorphism for all  $r$ .

*Proof of Lemma 2.* By definition,

$$\square_n = \{(x_1, \dots, x_n) \in \mathbb{R}^n : |x_i| \leq 1 \text{ for all } i = 1, \dots, n\} \quad (8)$$

$$\square_k = \{(x_1, \dots, x_k) \in \mathbb{R}^k : |x_i| \leq 1 \text{ for all } i = 1, \dots, k\}$$

$$Q = \{(x_1, \dots, x_n) \in \mathbb{R}^n : |x_i| \leq 1 + \varepsilon \text{ for } i = 1, \dots, k, \text{ and} \quad (9) \\ |x_i| \leq 1 - \varepsilon \text{ for } i = k + 1, \dots, n\}.$$

We identify  $\square_k$  with the cross section

$$\{(x_1, \dots, x_n) \in \square_n : x_{k+1} = \dots = x_n = 0\}.$$

By (8) and (9),

$$Q \setminus \square_n = \{(x_1, \dots, x_n) \in \mathbb{R}^n : |x_i| \leq 1 + \varepsilon \text{ for } i = 1, \dots, k \text{ and} \\ |x_i| \leq 1 - \varepsilon \text{ for } i = k + 1, \dots, n \text{ and} \\ |x_j| > 1 \text{ for some } j\}.$$

If  $j \geq k + 1$ , then  $|x_j| \leq 1 - \varepsilon < 1$ ; therefore  $|x_j| > 1$  is only possible when  $1 \leq j \leq k$ . Thus

$$Q \setminus \square_n = \left\{ (x_1, \dots, x_n) \in \mathbb{R}^n : \begin{array}{l} |x_i| \leq 1 + \varepsilon \text{ for } i = 1, \dots, k \\ |x_i| \leq 1 - \varepsilon \text{ for } i = k + 1, \dots, n \\ |x_j| > 1 \text{ for some } j, 1 \leq j \leq k \end{array} \right\}. \quad (10)$$

Next we define a retraction of  $Q \setminus \square_n$  onto  $\square_k(1 + \varepsilon) \setminus \square_k$ . For each  $t \in [0, 1]$ , set  $f_t(x_1, \dots, x_n) = (x_1, \dots, x_k, tx_{k+1}, \dots, tx_n)$ . If  $|x_i| \leq 1 - \varepsilon$  then  $|tx_i| \leq |x_i| \leq$

$1 - \varepsilon$ , so  $f_t(Q \setminus \square_n) \subset Q \setminus \square_n$  for all  $t$ . Note also that  $f_1$  is the identity map and  $f_0(x_1, \dots, x_n) = (x_1, \dots, x_k, 0, \dots, 0)$  lies in the set

$$\square_k(1 + \varepsilon) \setminus \square_k = \{(x_1, \dots, x_k, 0, \dots, 0) \quad : \quad |x_i| \leq 1 + \varepsilon \text{ for } i = 1, \dots, k \text{ and} \\ |x_j| > 1 \text{ for some } j = 1, \dots, k\}.$$

This proves that  $f_0$  is homotopic to the identity map and is a retraction. It follows that  $Q \setminus \square_n$  is homotopic to  $\square_k(1 + \varepsilon) \setminus \square_k$ .

On the other hand, we can show that  $\square_k(1 + \varepsilon) \setminus \square_k$  retracts onto  $\partial \square_k$ . For any  $x = (x_1, \dots, x_n)$ , let  $m(x) = \{|x_i| : 1 \leq i \leq n\}$ . If  $x \neq 0$  then  $m(x) \neq 0$ , so  $p(x) = \frac{1}{m(x)} \cdot x$  is well-defined. In particular,  $p$  is defined on  $\square_k(1 + \varepsilon) \setminus \square_k$  and maps this set onto  $\partial \square_k$ . The function  $p_t(x) := (\frac{1}{m(x)})^t \cdot x$ , for  $t \in [0, 1]$ , gives a homotopy from  $p_0 = id$  to  $p = p_1$ . This proves that  $p$  is a retraction.

In conclusion,  $p \circ f_0$  maps  $Q \setminus \square_n$  onto  $\partial \square_k$  and is a homotopy equivalence. As  $\partial \square_k$  is homotopy equivalent to  $S^{k-1}$ , the lemma is proved.  $\square$

We will need a preferred generator for the proof of Lemma 4 below, so let us specify one now. By Proposition 2 and Lemma 2,  $H_{k-1}(Q \setminus \square_n) \cong \mathbb{Z}$  has two possible generators. First, observe that  $\partial \square_k$  has a natural decomposition as a formal sum of  $(k-1)$ -simplices, and that  $\partial(\partial \square_k) = 0$ . It follows that  $\partial \square_k$  represents an element in  $\mathcal{Z}_{k-1}(\partial \square_k)$ . In fact, the homology class represented by  $\partial \square_k$  generates the group  $H_{k-1}(\partial \square_k) \cong \mathbb{Z}$ . We denote this class by  $[\partial \square_k]$ . Next let  $i_1 : \partial \square_k \hookrightarrow Q \setminus \square_n$  denote the natural inclusion. Note that  $i_1$  is a homotopy inverse to  $p \circ f_0$ .

**Definition 6.** The class  $[i_1] = (i_1)_*[\partial \square_k]$  will be our preferred generator for  $H_{k-1}(Q \setminus \square_n)$ .

*Proof of Lemma 3.* By definition,

$$\partial_X \square_n = \{(x_1, \dots, x_n) \in \square_n : |x_j| = 1 \text{ for some } j = 1, \dots, k\}. \quad (11)$$

For  $t \in [0, 1]$ , define  $g_t : \partial_X \square_n \rightarrow \partial_X \square_n$  by

$$g_t(x_1, \dots, x_n) = (x_1, \dots, x_k, tx_{k+1}, \dots, tx_n).$$

Note that if  $|x_i| \leq 1$ , then  $|tx_i| \leq |x_i| \leq 1$ , so indeed  $g_t(\partial_X \square_n) \subset \partial_X \square_n$ . As in the proof of Lemma 2,  $g_1$  is the identity map and  $g_0(x_1, \dots, x_n) = (x_1, \dots, x_k, 0, \dots, 0) \in \partial \square_k$ , which proves that  $g_0 \simeq g_1$ . Thus  $g_0$  is a retraction of  $\partial_X \square_n$  onto  $\partial \square_k$ .

On the other hand,  $\partial \square_k \simeq \partial D^k = S^{k-1}$ , so we have proved

$$\partial_X \square_n \simeq \partial \square_k \simeq S^{k-1},$$

as required.

*Remark.* Let  $i_2$  denote the natural inclusion of  $\partial \square_k$  into  $\partial_X \square_n$ , and note that  $i_2$  is a homotopy inverse to  $g_0$ . We will take  $[i_2] = (i_2)_*[\partial \square_k]$  to be the preferred generator of  $H_{k-1}(\partial_X \square_n) \cong \mathbb{Z}$ .

*Proof of Lemma 4.* Let  $\varphi$  be a diffeomorphism satisfying  $(n, k)$ -ICP, and  $\phi$  be the associated form of  $\varphi$  in standardized co-ordinates that satisfies the Standardized  $(n, k)$ -ICP. Then  $\phi$  maps  $\partial_X \square_n$  into the set  $Q \setminus \square_n$ . We claim that

i)  $\phi|_{\partial_X \square_n}$  is homotopic to  $h|_{\partial_X \square_n}$ , where  $h$  is the ‘‘hyperbolic map’’ defined by

$$h(x_1, \dots, x_n) = ((1 + \varepsilon)x_1, \dots, (1 + \varepsilon)x_k, (1 - \varepsilon)x_{k+1}, \dots, (1 - \varepsilon)x_n); \quad (12)$$

ii)  $h|_{\partial_X \square_n} : \partial_X \square_n \rightarrow Q \setminus \square_n$  induces an isomorphism in homology.

We begin by proving (i). For each  $j = 1, \dots, k$ , let

$$Q_{j+} = Q \cap \{x_j > +1\}, \quad Q_{j-} = Q \cap \{x_j < -1\}.$$

Thus, for example,  $Q_{j+}$  consists of all points  $(x_1, \dots, x_n)$  such that  $|x_i| \leq 1 + \varepsilon$  for  $1 \leq i \leq k$ ,  $|x_i| \leq 1 - \varepsilon$  for  $k+1 \leq i \leq n$ , and  $x_j > +1$ .

Each of these sets is a product of intervals; namely,

$$Q_{j+} = [-1 - \varepsilon, 1 + \varepsilon]^{k-1} \times (1, 1 + \varepsilon] \times [-1 + \varepsilon, 1 - \varepsilon]^{n-k}$$

$$Q_{j-} = [-1 - \varepsilon, 1 + \varepsilon]^{k-1} \times [-1 - \varepsilon, -1] \times [-1 + \varepsilon, 1 - \varepsilon]^{n-k}.$$

Therefore  $Q_{j\pm}$  is fully contractible.

By the Standardized  $(n, k)$ -ICP,  $\phi$  maps the set

$$B_{j\pm} := \partial_X \square_n \cap \{x_j = \pm 1\}$$

into  $Q_{j\pm}$ . The hyperbolic map  $h$  defined by (12) also maps  $B_{j\pm}$  into  $Q_{j\pm}$ . As  $Q_{j\pm}$  is contractible, the two maps

$$\phi|_{B_{j\pm}} : B_{j\pm} \rightarrow Q_{j\pm} \text{ and } h|_{B_{j\pm}} : B_{j\pm} \rightarrow Q_{j\pm}$$

must be homotopic. Let  $H_{j\pm} : B_{j\pm} \times I \rightarrow Q_{j\pm}$  be a homotopy between them with  $H_{j\pm}(\cdot, 0) = \phi$  and  $H_{j\pm}(\cdot, 1) = h$ .

For any  $j' \neq j$ ,  $H_{j\pm}$  maps the ‘‘overlap’’  $(B_{j\pm} \cap B_{j'\pm}) \times I$  into  $Q_{j\pm} \cap Q_{j'\pm}$ . The same is true for  $H_{j'\pm}$ . But the intersection of  $Q_{j\pm}$  and  $Q_{j'\pm}$  is also a product of intervals, hence contractible. This implies that we can choose the homotopy maps  $H_{j\pm}$  in such a way that they agree on overlaps; *i.e.*, if  $x \in B_{j\pm} \cap B_{j'\pm}$  and  $t \in I$ , then  $H_{j\pm}(x, t) = H_{j'\pm}(x, t)$ .

Now  $\partial_X \square_n$  equals the union, over all  $j = 1, \dots, k$ , of  $B_{j+} \cup B_{j-}$ . Thus, given any  $(x, t) \in \partial_X \square_n \times I$ , we can find  $j$  such that  $x \in B_{j+}$  or  $B_{j-}$ . We therefore construct a homotopy  $H : \partial\Gamma \times I \rightarrow Q \setminus \square_n$  from  $\phi = H(\cdot, 0)$  to  $h = H(\cdot, 1)$  by patching together the various maps  $H_{j\pm}$ . To be precise, if  $(x, t) \in \partial\Gamma \times I$ , then choose  $j \in \{1, \dots, k\}$  such that  $x$  lies in  $B_{j+}$  or  $B_{j-}$ . Suppose for example that  $x \in B_{j+}$ . Then we define  $H(x, t) := H_{j+}(x, t)$ . If  $x \in B_{j'\pm}$  for some other  $j' \neq j$  in the set  $\{1, \dots, k\}$ , then  $H_{j'\pm}(x, t) = H_{j+}(x, t)$  by the remarks of the preceding paragraph. Thus  $H$  is a well-defined map, and (i) is proved.

To prove (ii), we will show that  $h_*$  maps  $[i_2]$  to  $[i_1]$ . Thus we need to show that  $h_*(i_2)_*[\partial \square_k] = (i_1)_*[\partial \square_k]$ . Since  $i_1$  and  $p \circ f_0$  are homotopy inverses, it is equivalent to show that

$$(p \circ f_0)_* h_*(i_2)_*[\partial \square_k] = [\partial \square_k]. \quad (13)$$

Now consider  $p \circ f_0 \circ h \circ i_2$ , which maps  $\partial \square_k$  to itself. It is simple to check that for any  $(x_1, \dots, x_k, 0, \dots, 0) \in \partial \square_k$ ,

$$\begin{aligned} & p \circ f_0 \circ h \circ i_2(x_1, \dots, x_k, 0, \dots, 0) \\ &= p \circ f_0((1 + \varepsilon)x_1, \dots, (1 + \varepsilon)x_k, 0, \dots, 0) \\ &= p((1 + \varepsilon)x_1, \dots, (1 + \varepsilon)x_k, 0, \dots, 0) \\ &= \frac{1}{1 + \varepsilon} \cdot ((1 + \varepsilon)x_1, \dots, (1 + \varepsilon)x_k, 0, \dots, 0). \end{aligned}$$

Thus  $p \circ f_0 \circ h \circ i_2|_{\partial \square_k}$  equals the identity map. But the identity map induces the identity map in homology, so we have proved (13).

*Proof of Lemma 1.* By transversality,  $\Gamma' = \phi(\Gamma) \cap \square_n$  is a  $k$ -manifold with boundary, and its boundary equals  $\phi(\Gamma) \cap \partial \square_n$ . To see that  $\Gamma' \neq \emptyset$ , we use Lemma 4. For suppose that  $\phi(\Gamma) \cap \square_n = \emptyset$ . Then  $\phi(\Gamma) \subset Q \setminus \square_n$  and  $\partial(\phi(\Gamma)) = \phi(\partial\Gamma)$ , which

is to say that  $\phi(\partial\Gamma)$  is a boundary element in  $Q \setminus \square_n$ , *i.e.*, it represents an element in  $\mathcal{B}_{k-1}(Q \setminus \square_n)$ . By the definition of singular homology (7),  $[\phi(\partial\Gamma)] = 0$  in  $H_{k-1}(Q \setminus \square_n)$ . On the other hand,  $[\phi(\partial\Gamma)] = \phi_*[\partial\Gamma]$  because  $\phi$  is a diffeomorphism, and  $[\partial\Gamma] \neq 0$  by our wrap-around assumption. By Lemma 4,  $\phi_* : H_{k-1}(\partial_X \square_n) \rightarrow H_{k-1}(Q \setminus \square_n)$  is an isomorphism, meaning that  $\phi_*[\partial\Gamma] \neq 0$ , a contradiction. Thus our assumption on  $\phi(\Gamma) \cap \square_n$  must have been false. We conclude that  $\phi(\Gamma) \cap \square_n$  is non-empty. This proves (i).

By the Standardized  $(n, k)$ -ICP,  $\phi(\Gamma) \subset Q$  and so  $\phi(\Gamma) \cap \partial_C \square_n = \emptyset$ . This proves (ii).

Let  $\iota$  denote the inclusion  $\partial_X \square_n \hookrightarrow Q \setminus \square_n$ . We have

$$\iota \circ i_2 = i_1, \quad (14)$$

where  $i_1 : \partial \square_k \hookrightarrow Q \setminus \square_n$  and  $i_2 : \partial \square_k \hookrightarrow \partial_X \square_n$  denote, as before, the natural inclusion maps. Since  $[i_2]$  generates  $H_{k-1}(\partial_X \square_n) \cong \mathbb{Z}$ , there are uniquely determined integers  $d, d' \in \mathbb{Z}$  such that

$$[\partial\Gamma] = d \cdot [i_2] \quad \text{and} \quad [\partial\Gamma'] = d' \cdot [i_2].$$

Let  $A = \phi(\Gamma) \setminus \text{Int}(\square_n)$ . By our transversality assumption,  $A$  is a  $k$ -manifold with boundary, and its boundary equals

$$\phi(\partial\Gamma) - \partial\Gamma'. \quad (15)$$

This is to say that  $\partial A$  is the disjoint union of  $\phi(\partial\Gamma)$  and  $\partial\Gamma'$ , and that  $\partial\Gamma'$  is included with its orientation reversed (hence the minus sign in (15)). By the Theorem of Whitehead,  $A$  is triangulable, hence represents a class  $[A] \in C_k(Q \setminus \square_n)$  with the property that  $\partial[A] = [\phi(\partial\Gamma)] - [\iota(\partial\Gamma')]$ . By the definition of singular homology (7), we therefore have

$$[\phi(\partial\Gamma)] - [\iota(\partial\Gamma')] = 0 \quad (16)$$

in  $H_{k-1}(Q \setminus \square_n)$ .

Since  $\phi$  is a diffeomorphism,  $[\phi(\partial\Gamma)] = \phi_*[\partial\Gamma]$ . From the proof of Lemma 4 we also know that  $\phi \simeq h$  and  $h_*[i_2] = [i_1]$ . Therefore

$$[\phi(\partial\Gamma)] = \phi_*[\partial\Gamma] = h_*d \cdot [i_2] = d \cdot h_*[i_2] = d \cdot [i_1]. \quad (17)$$

On the other hand, by (14),

$$[\iota(\partial\Gamma')] = \iota_*[\partial\Gamma'] = \iota_*d' \cdot [i_2] = \iota_*(i_2)_*d' \cdot [\partial \square_k] = (i_1)_*d' \cdot [\partial \square_k] = d' \cdot [i_1]. \quad (18)$$

Combining (16), (17) and (18), we find that  $d[i_1] = d'[i_1]$ . Since  $[i_1]$  generates  $H_{k-1}(Q \setminus \square_n)$ , it follows that  $d' = d$ . Thus  $[\partial\Gamma'] = [\partial\Gamma]$ , and we are done.

*Acknowledgements* This research was conducted while the first two authors were postdoctoral fellows at the Fields Institute for Research in Mathematical Sciences, located in Toronto, Canada. We thank Judy Kennedy and Jim Yorke for helpful discussions.

## REFERENCES.

- Anosov, D. V. (1967). Geodesic Flows and Closed Riemannian Manifolds with Negative Curvature. *Proc. Steklov Inst. Math* 90, 1.
- Beyn, W.-J. (1987). On Invariant Closed Curves for One-Step Methods. *Numer. Math.* 51, 103–122.
- Bowen, R. (1975).  $\omega$ -Limit Sets for Axiom A Diffeomorphisms. *Journal of Differential Equations* 18, 333.

- Grebogi, C., S. M. Hammel, J. A. Yorke, and T. Sauer (1990, 24 September). Shadowing of Physical Trajectories in Chaotic Dynamics: Containment and Refinement. *Physical Review Letters* 65(13), 1527–1530.
- Greenberg, H. (1981). *Algebraic Topology: A First Course*. Addison-Wesley.
- Guillemin, V. and A. Pollack (1974). *Differential Topology*. Prentice-Hall.
- Hammel, S. M., J. A. Yorke, and C. Grebogi (1987). Do Numerical Orbits of Chaotic Dynamical Processes Represent True Orbits? *Journal of Complexity* 3, 136–145.
- Hammel, S. M., J. A. Yorke, and C. Grebogi (1988). Numerical Orbits of Chaotic Dynamical Processes Represent True Orbits. *Bull. Am. Math. Soc.* 19, 465–470.
- Hayes, W. B. (2001). *Rigorous shadowing of numerical solutions of ordinary differential equations by containment*. Ph. D. thesis, Department of Computer Science, University of Toronto. Available on the web as <http://www.cs.toronto.edu/NA/reports.html#hayes-01-phd>.
- Hayes, W. B. and K. R. Jackson (2003). Rigorous shadowing of numerical solutions of ordinary differential equations by containment. *SIAM J. Numer. Anal.* 41:5, 1948–1973.
- Milnor, J. W. and J. D. Stasheff (1974). Characteristic classes. *Annals of Mathematics Studies* (76).
- Palmer, K. J. (1988). Exponential Dichotomies, the Shadowing Lemma and Transversal Homoclinic Points. In U. Kirchgraber and H. O. Walther (Eds.), *Dynamics Reported*, Volume 1. Wiley and Teubner.
- Smale, S. (1965). Diffeomorphisms with many periodic points. In S. Cairns (Ed.), *Differential and Combinatorial Topology*, pp. 63–80. Princeton Univ. Press.
- Smale, S. (1967). Differentiable dynamical systems. *Bull. Amer. Math. Soc.* 73, 747–817.

Received September 15, 2004; in revised February 15, 2005.

*E-mail address:* wayne@ics.uci.edu