
Convergence Rate of Block-Coordinate Maximization Burer-Monteiro Method for Solving Large SDPs*

Murat A. Erdogdu[†], Asuman Ozdaglar[‡], Pablo A. Parrilo[‡], N. Denizcan Vanli[‡]

[†]University of Toronto, Vector Institute, erdogdu@cs.toronto.edu

[‡]Massachusetts Institute of Technology, {asuman,parrilo,denizcan}@mit.edu

Abstract

Semidefinite programming (SDP) with equality constraints arise in many optimization and machine learning problems, such as Max-Cut, community detection and robust PCA. Although SDPs can be solved to arbitrary precision in polynomial time, generic convex solvers do not scale well with the dimension of the problem. In order to address this issue, Burer and Monteiro [BM03] proposed to reduce the dimension of the problem by appealing to a low-rank factorization, and solve the subsequent non-convex problem instead. It is well-understood that the resulting non-convex problem acts as a reliable surrogate to the original SDP, and can be efficiently solved using the block-coordinate maximization method. Despite its simplicity, remarkable success, and wide use in practice, the theoretical understanding of the convergence of this method is limited. We prove that the block-coordinate maximization algorithm applied to the non-convex Burer-Monteiro approach enjoys a global sublinear rate without any assumptions on the problem, and a local linear convergence rate despite no local maxima is locally strongly concave. We illustrate our results through examples and numerical experiments.

1 Introduction

A variety of problems in statistical estimation and machine learning require solving a combinatorial optimization problem, which are often intractable [VB96]. Semidefinite programs (SDP) are commonly used as convex relaxations for these problems, providing efficient algorithms with approximate optimality [Par03]. A typically used SDP is

$$\begin{aligned} & \text{maximize} && \langle A, X \rangle && \text{(CVX)} \\ & \text{subject to} && X_{ii} = 1, \text{ for } i \in [n], \\ & && X \succeq 0, \end{aligned}$$

where $A, X \in \mathbb{R}^{n \times n}$ and $[n] = \{1, 2, \dots, n\}$. This problem appears as a convex relaxation to the celebrated Max-Cut problem [GW95], graphical model inference [EDM17], community detection problems [BBV16], and group synchronization [MMMO17].

Although SDPs serve as reliable relaxations to many combinatorial problems, the resulting convex problem is still computationally challenging. Interior point methods can solve SDPs to arbitrary accuracy in polynomial-time, but they do not scale well with the problem dimension n . A popular approach to remedy these limitations is to introduce a low-rank factorization $X = \sigma\sigma^\top$, where $\sigma \in \mathbb{R}^{n \times r}$ with r denoting the rank. This reformulation removes the positive semidefinite cone constraint in (CVX) since $X = \sigma\sigma^\top$ is guaranteed to be a positive semidefinite matrix, and choosing $r \ll n$ provides computational efficiency as well as storage benefits. This method is often referred to

*This work is supported by DARPA Lagrange program.

as Burer-Monteiro approach [BM03]. Denoting i -th row of σ by σ_i , i.e., $\sigma = [\sigma_1, \sigma_2, \dots, \sigma_n]^\top$, the resulting non-convex problem can be written as follows

$$\begin{aligned} & \text{maximize} && \langle A, \sigma \sigma^\top \rangle && \text{(Non-CVX)} \\ & \text{subject to} && \|\sigma_i\| = 1, \text{ for } i \in [n], \end{aligned}$$

where the non-convexity comes from the separable submanifold constraints $\|\sigma_i\| = 1$. In the original Burer-Monteiro approach [BM03], the authors propose to use an augmented Lagrangian method for a general form SDP. However, it has been recently observed that feasible methods (such as block-coordinate maximization [JMRT16, WCK17], Riemannian gradient [JMRT16, MMMO17] and Riemannian trust-region methods [ABG07, JBAS10, BVB16]) provide empirically faster rates since feasibility can be efficiently guaranteed via projection onto the Cartesian product of spheres. Despite many empirical evidence [JMRT16, MMMO17, WCK17], not much is known on the convergence of these feasible methods (except the Riemannian trust-region method, for which a sublinear convergence rate is shown in [BVB16] and a local superlinear convergence is shown in [ABG07] with no rate estimate). Among these methods, block-coordinate maximization and projected Riemannian gradient ascent are simpler to implement and have computational complexity of $\mathcal{O}(nr)$ and $\mathcal{O}(n^2r)$, respectively, whereas Riemannian trust-region requires the eigendecomposition of the dual variable (which is usually computed iteratively using the power method, whose each iteration requires $\mathcal{O}(n^2)$ arithmetic operations) and the ascent step requires an additional $\mathcal{O}(n^2r)$ complexity. Furthermore, block-coordinate maximization does not have any step size or tuning parameters, unlike projected Riemannian gradient ascent and Riemannian trust-region methods. Empirical studies further consolidate the use of block-coordinate maximization by presenting excellent results on many problems with often linear convergence. In this paper, we provide the first local and global convergence rate guarantees for the block-coordinate maximization method (applied to Burer-Monteiro approach) in the literature, which are consistent with empirical performance of the algorithm. Our contributions can be summarized as follows:

- We establish the global sublinear convergence of the block-coordinate maximization algorithm applied to the non-convex Burer-Monteiro approach without any assumptions on A .
- We show that the block-coordinate maximization algorithm enjoys a linear rate around a local neighborhood of any local maxima, even though no local maxima is locally strongly concave.
- Our complexity estimates yield an optimal sampling scheme of update blocks to tighten the sublinear and linear rates.
- We validate our theoretical results via numerical examples and compare the performance of the block-coordinate maximization algorithm with respect to various manifold optimization methods to emphasize its performance.

1.1 Related Work

There are numerous papers that analyze the landscape of (Non-CVX). In particular, it is known that (CVX) admits a maxima of rank at most $r \leq n(n+1)/2$ [Bar95, Pat98]. Using this observation, it has been shown in [BM03, BM05, JBAS10] that when $r \geq \sqrt{2n}$, if σ is a rank deficient second-order stationary point, then σ is a global maxima for (Non-CVX) and $X = \sigma \sigma^\top$ is a global maxima for (CVX). The recent paper [BVB18] showed that when $r \geq \sqrt{2n}$, for almost all A , every σ that is a first-order stationary point is rank deficient. For arbitrary rank r , [Mon16] showed that all local maxima are within a $n \|A\|_2 / \sqrt{r}$ gap from the (CVX) optimum, and [MMMO17] showed that any ε -approximate concave point is within a $\text{Rg}(\text{Non-CVX})/(r-1) + n\varepsilon/2$ gap from the (CVX) optimum, where $\text{Rg}(\text{Non-CVX})$ is the range of the problem (Non-CVX).

[JMRT16] presented that when applied to solve (Non-CVX), projected Riemannian gradient ascent and block-coordinate maximization methods provide excellent numerical results, yet no convergence guarantee is provided. Similar experimental results are also observed in [WCK17] for the block-coordinate maximization algorithm and [MMMO17] for the projected Riemannian gradient ascent algorithm. In [BAC16], the authors provided a global sublinear convergence rate for the Riemannian trust-region method for general non-convex problems and these results have been used in [BVB16, MMMO17] for the non-convex Burer-Monteiro approach. Augmented Lagrangian methods have been proposed to solve (Non-CVX) as well [BM03, BM05], however these methods do not benefit from separability of the manifold constraints, and hence are usually slower [BVB18].

1.2 Notations and Preliminaries

Throughout the paper, all vectors are column vectors. The superscripts are used to denote iteration counters, i.e., σ^k denotes the value of σ at iteration k . For a vector g , $\|g\|$ represents its Euclidean norm. For a matrix A , A_{ij} represents its entry at the i -th row and j -th column, $\|A\|_F$ represents its Frobenius norm, $\|A\|_1 = \max_{1 \leq j \leq n} \sum_{i=1}^n |A_{ij}|$ represents its 1-norm, and $\|A\|_{1,1} = \sum_{i,j=1}^n |A_{ij}|$ represents its $L_{1,1}$ -norm. For a function h , ∇h and $\text{grad}h$ represent its Euclidean and Riemannian gradient, respectively. Similarly, $\nabla^2 h$ and $\text{Hess}h$ represent its Euclidean and Riemannian Hessian, respectively. We let \mathbb{S}^{m-1} denote the unit sphere in \mathbb{R}^m .

Without loss of generality, we assume that A is symmetric and $A_{ii} = 0$, for all $i \in [n]$. Indeed, if A is not a symmetric matrix, then we can replace A by $(A + A^\top)/2$, which is a symmetric matrix, and the objective value (**Non-CVX**) remains the same for all $\sigma \in \mathbb{R}^{n \times r}$ since $\sigma\sigma^\top$ is symmetric. Similarly, replacing the diagonal entries of A by zeros decreases the objective value by the constant $\text{Tr}(A)$, for all $\sigma \in \mathbb{R}^{n \times r}$ since the diagonal entries of $\sigma\sigma^\top$ are all equal to 1.

The rest of the paper is organized as follows. In Section 2, we present the algorithm, discuss its complexity and compare it to the other feasible methods. In Section 3, we prove the global sublinear convergence of the algorithm and provide rate estimates. In Section 4, we show that the algorithm enjoys a local linear convergence rate and provide rate estimates. We perform numerical experiments to validate our theoretical results in Section 5 and conclude the paper in Section 6.

2 Block-Coordinate Maximization (BCM) Algorithm

In this section, we discuss the update rule and computational complexity of the BCM algorithm. Given the current iterate σ^k , the BCM algorithm chooses a block σ_{i_k} and maximizes the objective

$$f(\sigma^k) = \sum_{i=1}^n \langle \sigma_i^k, g_i^k \rangle, \quad \text{where } g_i^k := \sum_{j \neq i} A_{ij} \sigma_j^k,$$

over $\sigma_{i_k} \in \mathbb{S}^{r-1}$. More formally, we can write the update rule of the algorithm as follows

$$\begin{aligned} \sigma_{i_k}^{k+1} &= \arg \max_{\|\sigma\|=1} f(\sigma_1^k, \dots, \sigma_{i_k-1}^k, \sigma, \sigma_{i_k+1}^k, \dots, \sigma_n^k) \\ &= \arg \max_{\|\sigma\|=1} \langle \sigma, g_{i_k}^k \rangle + \sum_{i \neq i_k} \sum_{j \neq i} A_{ij} \langle \sigma_i^k, \sigma_j^k \rangle \\ &= \arg \max_{\|\sigma\|=1} 2 \langle \sigma, g_{i_k}^k \rangle + \sum_{i \neq i_k} \sum_{j \neq i, i_k} A_{ij} \langle \sigma_i^k, \sigma_j^k \rangle \\ &= \arg \max_{\|\sigma\|=1} \langle \sigma, g_{i_k}^k \rangle = \frac{g_{i_k}^k}{\|g_{i_k}^k\|}, \end{aligned} \tag{1}$$

with the convention that $\sigma_{i_k}^{k+1}$ can be chosen arbitrarily when $\|g_{i_k}^k\| = 0$, and where the third equality follows since A is symmetric. Although i_k can be chosen arbitrarily, we focus on random selection in this paper. In particular, we consider two randomization schemes:

- $i_k = i$ with probability $p_i = 1/n$, which we call as uniform sampling,
- $i_k = i$ with probability $p_i = \|g_i^k\| / \sum_{j=1}^n \|g_j^k\|$, which we call as importance sampling.

The BCM algorithm with uniform sampling can be implemented in $\mathcal{O}(nr)$ time and space complexity since it only needs to save σ (which is of size nr) and after i_k is chosen $g_{i_k}^k$ can be computed in $2(n-1)r$ floating point operations. On the other hand, the BCM algorithm with importance sampling requires $\|g_i^k\|$, for all $i \in [n]$, to be able to assign p_i 's. Although each g_i^k can be computed at the beginning of each iteration, this would require $\mathcal{O}(n^2r)$ floating point operations. Instead, a smarter way of implementation would be to keep both σ and g_i 's in the memory (which requires $2nr$ space) and update $\sigma_{i_k}^k$ and g_i^k , for all $i \neq i_k$ as presented in Algorithm 1, which yields $2(n-1)r$ floating point operations to update g_i 's similar to the uniform sampling case. Therefore, the time and space complexity of both variants are $\mathcal{O}(nr)$ for dense A (i.e., when no structure is available

Algorithm 1 Block-Coordinate Maximization (BCM)

Initialize $\sigma^0 \in \mathbb{R}^{n \times r}$ and calculate $g_i^0 = \sum_{j \neq i} A_{ij} \sigma_j^0$, for all $i \in [n]$.

for $k = 0, 1, 2, \dots$ **do**

Choose block $i_k = i$, with probability p_i , for all $i \in [n]$,

$$\sigma_{i_k}^{k+1} \leftarrow \frac{g_{i_k}^k}{\|g_{i_k}^k\|},$$

$$g_i^{k+1} \leftarrow g_i^k - A_{ii_k} \sigma_{i_k}^k + A_{ii_k} \sigma_{i_k}^{k+1}, \text{ for all } i \neq i_k.$$

end for

on A). However, in many SDP applications (such as Max-Cut and graphical model inference), A is induced by a graph. Therefore, the computational cost of the BCM algorithm can be reduced to $\mathcal{O}(dr)$, where d is the maximum degree of the graph that induces A . In comparison, per iteration computational complexity of the projected Riemannian gradient ascent algorithm is $\mathcal{O}(n^2r)$ for dense A and $\mathcal{O}(d^2r)$ for sparse A . The situation is even worse for Riemannian trust-region algorithm since it requires to perform power method to solve the trust-region subproblem to find an approximate update direction. Hence, per iteration complexity of the BCM algorithm is much smaller than the other feasible methods. Furthermore, the BCM algorithm has structural advantages over projected Riemannian gradient ascent as well. In particular, the fixed points of the BCM algorithm are of the form $\sigma_i = g_i / \|g_i\|$, whereas any first-order stationary point, i.e., σ 's that satisfy $\sigma_i = \pm g_i / \|g_i\|$ (as we will show in (8)), is a fixed point of the projected Riemannian gradient ascent algorithm. We can observe that for a first-order stationary σ , if there exists a block ℓ such that $\sigma_\ell = -g_\ell / \|g_\ell\|$ holds, then we can increase $f(\sigma)$ by replacing σ_ℓ with any other unit norm vector since

$$\begin{aligned} f(\sigma_1, \dots, \sigma_\ell, \dots, \sigma_n) &= 2\langle \sigma_\ell, g_\ell \rangle + \sum_{i \neq \ell} \sum_{j \neq \ell, i} A_{ij} \langle \sigma_i, \sigma_j \rangle \\ &< 2\langle \bar{\sigma}_\ell, g_\ell \rangle + \sum_{i \neq \ell} \sum_{j \neq \ell, i} A_{ij} \langle \sigma_i, \sigma_j \rangle = f(\sigma_1, \dots, \bar{\sigma}_\ell, \dots, \sigma_n), \end{aligned}$$

for any $\bar{\sigma}_\ell \neq \sigma_\ell$ due to the Cauchy-Schwarz inequality. Taking the limit as $\bar{\sigma}_\ell \rightarrow \sigma_\ell$, we can observe that σ cannot be a local maxima. Therefore, we conclude that all local maxima satisfy $\sigma_i = g_i / \|g_i\|$, and hence the first-order stationary points that are not fixed points of BCM are either local minima or saddle. This intrinsically rules out convergence to exponentially many (in n) saddle points and local minima. Furthermore, projected Riemannian gradient ascent requires line-search to find a step size that yields an ascent at each step. On the other hand, as we show in Lemma 3.1, the BCM algorithm is guaranteed to make an ascent at each iteration without any parameter tuning.

3 Global Sublinear Convergence Rate

In this section, we prove that the BCM algorithm attains a global sublinear convergence and provide rate estimates. To this end, we first introduce the following ascent lemma, which shows that the sequence of function values $\{f(\sigma^k)\}_{k \geq 0}$ is nondecreasing.

Lemma 3.1. *Each iteration of the BCM algorithm yields the following ascent on the function value:*

$$f(\sigma^{k+1}) - f(\sigma^k) = 2(\|g_{i_k}^k\| - \langle \sigma_{i_k}^k, g_{i_k}^k \rangle) \geq 0.$$

We emphasize that such an ascent lemma does not necessarily hold for general non-convex functions and algorithms. In particular, in order to guarantee ascent condition, it is often required to use line-search techniques for choosing the step size of first-order methods (e.g., the gradient ascent algorithm) [SU15]. On the other hand, the BCM algorithm does not require any parameter tuning and still enjoys the ascent guarantee in Lemma 3.1. This lemma holds a basis for the following theorem, in which we consider the BCM algorithm with importance sampling and show that its expected functional ascent can be related to the expected norm of the Riemannian gradient of the function evaluated at the current iterate. Hence, it is guaranteed that the BCM algorithm returns a solution with arbitrarily small Riemannian gradient as we highlight in the following theorem.

Theorem 3.2. *Let $f^* = \max_{\|\sigma_i\|=1, \forall i \in [n]} f(\sigma)$. Then, in at most $K \geq \lceil (\|A\|_{1,1} (f^* - f(\sigma^0))) / \epsilon \rceil$ iterations, BCM with importance sampling is guaranteed to return a solution σ^k , for some $k \in [K-1]$,*




Figure 1: Convergence of the BCM algorithm, where the entries of A are drawn from a normal distribution and $n = 500$. The local convergence can be observed to be linear.

satisfying $\mathbb{E} \|\text{grad} f(\sigma^k)\|_{\text{F}}^2 \leq \epsilon$. Equivalently, for any $K \geq 1$, BCM with importance sampling yields the following guarantee

$$\min_{k \in [K-1]} \mathbb{E} \|\text{grad} f(\sigma^k)\|_{\text{F}}^2 \leq \frac{\|A\|_{1,1} (f^* - f(\sigma^0))}{K}. \quad (2)$$

Using a similar approach to Theorem 3.2, we show in the following corollary that the BCM algorithm with uniform sampling attains a similar sublinear convergence rate. However, comparing the rate guarantees in (2) and (3), we can observe that the BCM algorithm with importance sampling enjoys a tighter convergence rate compared to the BCM algorithm with uniform sampling, since $\|A\|_{1,1} \leq n \|A\|_1$, for all $A \in \mathbb{R}^{n \times n}$.

Corollary 3.3. *Let $f^* = \max_{\|\sigma_i\|=1, \forall i \in [n]} f(\sigma)$. Then, in at most $K \geq \lceil (n \|A\|_1 (f^* - f(\sigma^0))) / \epsilon \rceil$ iterations, BCM with uniform sampling is guaranteed to return a solution σ^k , for some $k \in [K-1]$, satisfying $\mathbb{E} \|\text{grad} f(\sigma^k)\|_{\text{F}}^2 \leq \epsilon$. Equivalently, for any $K \geq 1$, BCM with uniform sampling yields the following guarantee*

$$\min_{k \in [K-1]} \mathbb{E} \|\text{grad} f(\sigma^k)\|_{\text{F}}^2 \leq \frac{n \|A\|_1 (f^* - f(\sigma^0))}{K}. \quad (3)$$

4 Local Linear Convergence Rate

Although the BCM algorithm enjoys the sublinear convergence rates presented in Section 3, it is numerically observed that the rate of convergence is linear when σ^k is close to a local maxima [JMRT16, WCK17]. A similar conclusion can be made by Figure 1 as well, which illustrates local linear convergence of BCM. In this section, we investigate this behavior and prove that indeed BCM attains a linear convergence rate around a local maxima. In order to prove this result, we require certain tools from manifold optimization [AMS07]. We define the following submanifold of full-rank² matrices $\mathbb{R}_*^{n \times r}$ that corresponds to the Riemannian geometry induced by the constraints of the problem (Non-CVX) in the Euclidean space:

$$\mathcal{M}_r := \{\sigma = (\sigma_1, \dots, \sigma_n)^\top \in \mathbb{R}_*^{n \times r} : \|\sigma_i\| = 1, \forall i \in [n]\}.$$

This manifold represents the Cartesian product of n unit spheres in \mathbb{R}^r . For any given point $\sigma \in \mathcal{M}_r$, its tangent space can be found (by taking the differential of the equality constraints) as follows

$$T_\sigma \mathcal{M}_r := \{u = (u_1, \dots, u_n)^\top \in \mathbb{R}_*^{n \times r} : \langle u_i, \sigma_i \rangle = 0, \forall i \in [n]\}.$$

Using these definitions, the geodesics $t \mapsto \sigma(t)$ (i.e., curves of shortest path with zero acceleration) can be expressed as a function of $\sigma = \sigma(0) \in \mathcal{M}_r$ and $u \in T_\sigma \mathcal{M}_r$ as follows

$$\sigma_i(t) = \sigma_i \cos(\|u_i\| t) + \frac{u_i}{\|u_i\|} \sin(\|u_i\| t). \quad (4)$$

²We consider full-rank matrices to deal with differentiable manifolds, which is standard, e.g., see [JBAS10].

We refer to Section 5.4 of [AMS07] for a more detailed treatment of this topic. The above geodesic can be thought as the set of points on the manifold that are obtained by moving from $\sigma \in \mathcal{M}_r$ towards the direction pointed by $u \in T_\sigma \mathcal{M}_r$. Before understanding the landscape around a local maxima $\sigma \in \mathcal{M}_r$, we first make the following observation. Let $O(r) = \{Q \in \mathbb{R}^{r \times r} : Q^\top Q = QQ^\top = I\}$ denote the orthogonal group in dimension r . We can observe that $f(\sigma Q) = \langle A, \sigma Q Q^\top \sigma^\top \rangle = \langle A, \sigma \sigma^\top \rangle = f(\sigma)$, for any $Q \in O(r)$. Thus, every local maxima is flat in certain directions in $T_\sigma \mathcal{M}_r$. In order to characterize these directions, we define $\bar{\mathcal{M}}_r = \mathcal{M}_r / O(r)$ as the quotient of the manifold \mathcal{M} by the orthogonal group $O(r)$, which can be thought as the set of equivalence classes. We then consider the tangent space $T_\sigma \mathcal{M}_r$ and decompose it into two orthogonal subspaces: the vertical space $\mathcal{V}_\sigma \bar{\mathcal{M}}_r$ and the horizontal space $\mathcal{H}_\sigma \bar{\mathcal{M}}_r$. The vertical space $\mathcal{V}_\sigma \bar{\mathcal{M}}_r$ is the tangent space to equivalence classes, i.e.,

$$\mathcal{V}_\sigma \bar{\mathcal{M}}_r = \{\sigma Q : Q \in \mathbb{R}^{r \times r}, Q^\top = -Q\}.$$

This space contains the tangent vectors along which function value does not change and hence there is no curvature. The horizontal space $\mathcal{H}_\sigma \bar{\mathcal{M}}_r$ is the orthogonal complement of $\mathcal{V}_\sigma \bar{\mathcal{M}}_r$ in $T_\sigma \mathcal{M}_r$, i.e.,

$$\mathcal{H}_\sigma \bar{\mathcal{M}}_r = \{u \in T_\sigma \mathcal{M}_r : u^\top \sigma = \sigma^\top u\}.$$

In other words, $\mathcal{H}_\sigma \bar{\mathcal{M}}_r$ contains tangent vectors that do not rotate σ at all, which are the directions along which there is curvature. For a more detailed treatment of these definitions, we refer to Chapter 4 of [JBAS10], where similar equivalence class definitions are introduced to guarantee rotational invariance to design an algorithm, whereas our purpose here is to obtain convergence rate estimates.³ The main assumption we will use in proving the local linear convergence of the BCM algorithm is that along any direction in $\mathcal{H}_\sigma \bar{\mathcal{M}}_r$, $f(\sigma(t))$ has a negative curvature of at least $\mu > 0$. More formally, we make the following assumption.

Assumption 1. *Let σ be a local maxima of the problem (Non-CVX). Then, $\langle u, \text{Hess}f(\sigma)[u] \rangle \leq -\mu \|u\|_{\mathbb{F}}^2$ holds for all $u \in \mathcal{H}_\sigma \bar{\mathcal{M}}_r$.*

We emphasize that this assumption implies having isolated maximizers on the search space $\bar{\mathcal{M}}$, which is the assumption used in [JBAS10]. To make the above discussion and this assumption more vivid, we consider the following example.

Example: Consider the problem (Non-CVX), where A , σ , and u are defined as follows

$$A = \begin{bmatrix} 0 & \sqrt{2} & -1 \\ \sqrt{2} & 0 & \sqrt{2} \\ -1 & \sqrt{2} & 0 \end{bmatrix}, \quad \sigma = \begin{bmatrix} 1 & 0 \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ 0 & 1 \end{bmatrix}, \quad u = \begin{bmatrix} 0 & a \\ \frac{b}{\sqrt{2}} & \frac{-b}{\sqrt{2}} \\ c & 0 \end{bmatrix},$$

for some $a, b, c \in \mathbb{R}$. We can observe that σ is a local (global) maxima of the problem with $\sigma_i = g_i / \|g_i\|$, where $\|g_1\| = \|g_3\| = 1$ and $\|g_2\| = 2$. We can also see that any matrix in $T_\sigma \mathcal{M}_2$ is of the form u , for some $a, b, c \in \mathbb{R}$. Hence, letting $\Lambda := \text{diag}(\|g_1\|, \|g_2\|, \|g_3\|)$, we have

$$\frac{1}{2} \langle u, \text{Hess}f(\sigma)[u] \rangle = \langle u, (A - \Lambda)u \rangle = -(a+b)^2 - (c-b)^2,$$

which implies that $\langle u, \text{Hess}f(\sigma)[u] \rangle = 0$ if $b = c = -a$, and $\langle u, \text{Hess}f(\sigma)[u] \rangle < 0$ otherwise. Let \bar{u} denote the matrix u with entries $b = c = -a = 1$. Then, if we consider the geodesics $\sigma(t)$ defined by $\sigma \in \mathcal{M}_2$ and $\bar{u} \in T_\sigma \mathcal{M}_2$, i.e.,

$$\sigma_i(t) = \sigma_i \cos(\|\bar{u}_i\| t) + \frac{\bar{u}_i}{\|\bar{u}_i\|} \sin(\|\bar{u}_i\| t) = \sigma_i \cos(t) + \bar{u}_i \sin(t),$$

we can observe that $\sigma_i(t)$'s correspond to rotation of σ_i 's, for any $t \in \mathbb{R}$, i.e., $\sigma_i(t) = \sigma_i Q(t)$, for all i , for some rotation matrix $Q(t) \in O(2)$. Consequently, we have $\bar{u} \in \mathcal{F}_\sigma$ and by the above discussion we can conclude that $\mathcal{F}_\sigma = \text{span}(\bar{u})$. Hence, considering the directions in the tangent space $u \in T_\sigma \mathcal{M}_2$ that are orthogonal to \mathcal{F}_σ , we have

$$\begin{aligned} \max_{\substack{u \in T_\sigma \mathcal{M}_2 : \\ \|u\|_{\mathbb{F}}=1, \langle u, \bar{u} \rangle=0}} \frac{1}{2} \langle u, \text{Hess}(f(\sigma))[u] \rangle &= \max_{\substack{a^2+b^2+c^2=1, \\ -a+b+c=0}} -(a+b)^2 - (c-b)^2 \\ &= \max_{\substack{a^2+b^2+c^2=1, \\ -a+b+c=0}} -2 - b^2 + 2ac = -1, \end{aligned}$$

³An alternative approach for rotational invariance would be to consider adding additional constraints to (Non-CVX) such as imposing that σ is lower triangular, which defines a Cholesky manifold embedded in $\mathbb{R}^{n \times r}$ [GP07].

which is attained for $a = -c = \pm \frac{1}{\sqrt{2}}$ and $b = 0$. This example illustrates that Assumption 1 holds with constant $\mu = 2$ and although strong concavity does not hold for all $u \in T_\sigma \mathcal{M}_2$, it holds for a subset of matrices in the tangent space.

Before stating the main theorem on the local convergence rate of the BCM algorithm, we first present the following lemma, which states that the sequence $\{f(\sigma^k)\}_{k \geq 0}$ converges and the limit points of the sequence $\{\sigma^k\}_{k \geq 0}$ are the fixed points of the algorithm.

Lemma 4.1. *Let $\{\sigma^k\}_{k \geq 0}$ be the sequence generated by the BCM algorithm. Then, $\lim_{k \rightarrow \infty} f(\sigma^k)$ exists and every limit point $\bar{\sigma}$ of $\{\sigma^k\}_{k \geq 0}$ is a stationary point satisfying $\bar{\sigma}_i = \frac{\bar{g}_i}{\|\bar{g}_i\|}$.*

In the following theorem, we state the main linear convergence rate result for the BCM algorithm. An informal version of this theorem can be stated as follows. Suppose the BCM algorithm converges to a local maxima, for which Assumption 1 holds with some constant μ . Then, the algorithm attains a local linear convergence rate of $1 - \mu/(n \|A\|_{1,1})$ per iteration and $1 - \mu/\|A\|_{1,1}$ per cycle, approximately. In the formal statement of the theorem, we consider the case the sequence $\{\sigma^k\}_{k \geq 0}$ does not converge but instead has distinct limit points due to Lemma 4.1, where we emphasize that all limit points have the same function value due to Lemma 3.1.

Theorem 4.2. *Let $\bar{f} = \lim_{k \rightarrow \infty} f(\sigma^k)$, suppose Assumption 1 holds and assume that the limit points $\bar{\sigma}$ of the BCM algorithm are local maxima. Then, there exists an integer $K > 0$ such that the iterates generated by the BCM algorithm with importance sampling enjoy the following linear convergence rate*

$$\bar{f} - f(\sigma^{k+1}) \leq \left(1 - \frac{\mu}{n \|A\|_{1,1}} + \delta_K\right) (\bar{f} - f(\sigma^k)), \quad (5)$$

for any $k \geq K$, where δ_K is a constant that goes to 0 as $K \rightarrow \infty$.

Similar to the importance sampling case, we have the following linear convergence rate guarantee for the BCM algorithm with uniform sampling.

Corollary 4.3. *Let the conditions in Theorem 4.2 hold. Then, the iterates generated by the BCM algorithm with uniform sampling enjoy the same local linear convergence rate given in (5), with $\|A\|_{1,1}$ replaced by $n \|A\|_1$.*

5 Numerical Experiments

In this section, we compare the BCM algorithm with the projected Riemannian gradient ascent (PRGA) algorithm and the Riemannian trust-region (RTR) algorithm all applied to the Burer-Monteiro formulation in (Non-CVX). The RTR algorithm is implemented using the Manopt package [BMAS14] with the default options, while the gradient norm tolerance of the solution, i.e., $\|\nabla f(\sigma^k)\|_{\mathbb{F}} \leq \epsilon$, is set to $\epsilon = 10^{-5}$. The PRGA algorithm is implemented with a fixed step size of $5/\|A\|_1$, which provided the most consistent convergence throughout the experiments. In order to obtain a fair comparison between these first-order and second-order algorithms, we run the RTR algorithm to a desired gradient norm tolerance and count the elapsed time. We run the BCM and PRGA algorithms for the same amount of time (regardless of the number of iterations) and plot the Frobenius norms of the gradients and the suboptimality in function values with respect to the objective value each algorithm converges to, with respect to time elapsed. We emphasize that the BCM algorithm has a computational complexity of $\mathcal{O}(nr)$, whereas it is $\mathcal{O}(n^2r)$ for the PRGA and RTR algorithms with the caveat that RTR requires to run power method at each iteration, which costs $\mathcal{O}(n^2)$ arithmetic operations per inner loop. Therefore, one can expect given a fixed amount of time, BCM performs n times as many iterations as PRGA, while RTR performs less number of iterations than both.

In our experiments, we consider the case where A is randomly generated. In particular, we draw each entry of A from a normal distribution, symmetrize it, remove the diagonals and normalize it such that $\|A\|_{1,1} = 100$, for implementation convenience. As discussed in Section 1.2, these modifications do not change the landscape of the problem. In Figure 2, we plot the function suboptimality in the top row and the Frobenius norm of the Riemannian gradients in the bottom row. Each column corresponds to a separate problem. In the left column, we consider the case a relatively low-dimensional problem $n = 1000$ with low-rank $r = 20$. It can be observed that BCM rapidly decreases the function suboptimality and gradient norm, whereas RTR lags in the beginning and




Figure 2: Convergence of the BCM algorithm versus RTR and PRGA algorithms, where the entries of A are drawn from a normal distribution.

then attains a better convergence rate, which is expected since RTR converges locally superlinearly [ABG07]. On the other hand, PRGA is often better compared to RTR in a small time horizon, while its local convergence rate is similar to BCM's, which makes it fairly unfavorable compared to BCM and RTR. In the middle column in Figure 2, we consider the same dimension as the previous example, while setting $r = 45$, which satisfies $r \geq \sqrt{2n}$ and hence all stationary points are rank deficient with probability 1, which implies all local maxima are global [BVB18, JBAS10]. Changing r does not seem to affect the performance comparison between algorithms, while it is important to note that when r is significantly small (e.g., $r \leq 5$), it in general takes longer time for the first-order algorithms escape saddle points compared to RTR. In the right column in Figure 2, we consider a moderate size problem while keeping the rank same in comparison to the setting in the middle column. We observe that BCM and RTR perform much better than PRGA. Furthermore, BCM seems to obtain a better function suboptimality than RTR. Although we do not plot the comparison between the objective value that each algorithm attains in the limit, we observed (in all the experiments) that PRGA gets stuck in saddle points significantly more often than BCM and RTR. This is in accordance with the discussion we made earlier, where we noted that BCM escapes exponentially many saddles (in n) compared to PRGA by construction. On the other hand, the objective values BCM and RTR attain are usually close to one another, e.g., in the final example the objective value of the point returned by BCM is 5×10^{-4} higher than the objective value of the point returned by RTR, which is about the gap we observe in their performances and hence they both attain similar objective values in convergence. It can also be observed that as n increases, BCM tends to outperform RTR since the complexity of BCM scales better with n compared to RTR.

6 Conclusion

In this paper, we studied the non-convex Burer-Monteiro approach to solve large-scale SDPs. We considered to solve this non-convex problem using the block-coordinate maximization algorithm, which is significantly simpler to implement and computationally $\mathcal{O}(n)$ faster with respect to its alternatives such as the projected Riemannian gradient ascent and Riemannian trust-region methods. We proved that when the block-coordinates are chosen at random independently for each iteration, then the resulting algorithm attains a global sublinear convergence rate of $\mathcal{O}(1/\epsilon)$ to guarantee $\mathbb{E} \|\text{grad} f(\sigma^k)\|_F^2 \leq \epsilon$. We then analyzed the local behavior of the algorithm around a local maxima under a curvature assumption in a subset of the neighborhood of the local maxima. We showed that the block-coordinate maximization algorithm attains a local linear convergence rate of approximately $1 - \mu/(n \|A\|_{1,1})$ per iteration, which translates into an approximate rate of $1 - \mu/\|A\|_{1,1}$ per cycle, when $\mu/\|A\|_{1,1} \ll 1$. We presented the first precise rate estimates for the non-convex Burer-Monteiro approach in the literature to our knowledge. We validated our theoretical results through numerical experiments.

References

- [ABG07] P.-A. Absil, C.G. Baker, and K.A. Gallivan. Trust-region methods on riemannian manifolds. *Foundations of Computational Mathematics*, 7(3):303–330, Jul 2007.
- [AMS07] P.-A. Absil, R. Mahony, and R. Sepulchre. *Optimization Algorithms on Matrix Manifolds*. Princeton University Press, Princeton, NJ, USA, 2007.
- [BAC16] N. Boumal, P.-A. Absil, and C. Cartis. Global rates of convergence for nonconvex optimization on manifolds. *arXiv preprint arXiv:1605.08101*, 2016.
- [Bar95] A. I. Barvinok. Problems of distance geometry and convex properties of quadratic maps. *Discrete & Computational Geometry*, 13(2):189–202, 1995.
- [BBV16] A. S. Bandeira, N. Boumal, and V. Voroninski. On the low-rank approach for semidefinite programs arising in synchronization and community detection. *ArXiv:1602.04426*, 2016.
- [BM03] S. Burer and R. D. C. Monteiro. A nonlinear programming algorithm for solving semidefinite programs via low-rank factorization. *Mathematical Programming*, 95(2):329–357, 2003.
- [BM05] S. Burer and R. D. C. Monteiro. Local minima and convergence in low-rank semidefinite programming. *Mathematical Programming*, 103(3):427–444, Jul 2005.
- [BMAS14] N. Boumal, B. Mishra, P.-A. Absil, and R. Sepulchre. Manopt, a Matlab toolbox for optimization on manifolds. *Journal of Machine Learning Research*, 15:1455–1459, 2014.
- [BVB16] N. Boumal, V. Voroninski, and A. S. Bandeira. The non-convex Burer-Monteiro approach works on smooth semidefinite programs. In *Advances in Neural Information Processing Systems*, pages 2757–2765, 2016.
- [BVB18] N. Boumal, V. Voroninski, and A. S. Bandeira. Deterministic guarantees for Burer–Monteiro factorizations of smooth semidefinite programs. *arXiv preprint arXiv:1804.02008*, 2018.
- [EDM17] M. A. Erdogdu, Y. Deshpande, and A. Montanari. Inference in graphical models via semidefinite programming hierarchies. In *Advances in Neural Information Processing Systems*, pages 416–424, 2017.
- [GP07] I. Grubisic and R. Pietersz. Efficient rank reduction of correlation matrices. *Linear Algebra and its Applications*, 422(2):629 – 653, 2007.
- [GW95] M. X. Goemans and D. P. Williamson. Improved approximation algorithms for maximum cut and satisfiability problems using semidefinite programming. *Journal of the ACM (JACM)*, 42(6):1115–1145, 1995.
- [JBAS10] M. Journée, F. Bach, P.-A. Absil, and R. Sepulchre. Low-rank optimization on the cone of positive semidefinite matrices. *SIAM Journal on Optimization*, 20(5):2327–2351, 2010.
- [JMRT16] A. Javanmard, A. Montanari, and F. Ricci-Tersenghi. Phase transitions in semidefinite relaxations. *Proceedings of the National Academy of Sciences*, 113(16):E2218–E2223, 2016.
- [MMMO17] S. Mei, T. Misiakiewicz, A. Montanari, and R. I. Oliveira. Solving SDPs for synchronization and MaxCut problems via the Grothendieck inequality. *arXiv preprint arXiv:1703.08729*, 2017.
- [Mon16] A. Montanari. A Grothendieck-type inequality for local maxima. *arXiv preprint arXiv:1603.04064*, 2016.
- [Par03] P. A. Parrilo. Semidefinite programming relaxations for semialgebraic problems. *Mathematical Programming*, 96(2):293–320, May 2003.
- [Pat98] G. Pataki. On the rank of extreme matrices in semidefinite programs and the multiplicity of optimal eigenvalues. *Mathematics of operations research*, 23(2):339–358, 1998.
- [SU15] R. Schneider and A. Uschmajew. Convergence results for projected line-search methods on varieties of low-rank matrices via lojasiewicz inequality. *SIAM Journal on Optimization*, 25(1):622–646, 2015.
- [VB96] L. Vandenberghe and S. Boyd. Semidefinite programming. *SIAM Review*, 38(1):49–95, 1996.
- [WCK17] P.-W. Wang, W.-C. Chang, and J. Z. Kolter. The mixing method: coordinate descent for low-rank semidefinite programming. *arXiv preprint arXiv:1706.00476*, 2017.

A Calculation of Riemannian Gradient and Hessian

We define the following submanifold of $\mathbb{R}^{n \times r}$ that corresponds to the Riemannian geometry induced by the constraints of the problem (**Non-CVX**) in the Euclidean space:

$$\mathcal{M}_r := \{ \sigma = (\sigma_1, \dots, \sigma_n)^\top \in \mathbb{R}_*^{n \times r} : \|\sigma_i\| = 1, \forall i \in [n] \}, \quad (6)$$

This manifold represents the Cartesian product of n unit spheres in \mathbb{R}^r (which we denote by \mathbb{S}^{r-1}). For any given point $\sigma \in \mathcal{M}_r$, its tangent space can be found (by taking the derivatives of the sphere constraint) as follows

$$T_\sigma \mathcal{M}_r = \{ u = (u_1, \dots, u_n)^\top \in \mathbb{R}^{n \times r} : \langle u_i, \sigma_i \rangle = 0, \forall i \in [n] \}. \quad (7)$$

Before defining the Riemannian gradient and Hessian, we first let $P^\perp : \mathbb{R}^{n \times r} \rightarrow T_\sigma \mathcal{M}_r$ denote the projection operator from the Euclidean space to the tangent space of σ . When applied to a given matrix $w = (w_1, \dots, w_n)^\top \in \mathbb{R}^{n \times r}$, this projection operator yields

$$\begin{aligned} P^\perp(w) &= (P_1^\perp(w_1), \dots, P_n^\perp(w_n))^\top \\ &= (w_1 - \langle \sigma_1, w_1 \rangle \sigma_1, \dots, w_n - \langle \sigma_n, w_n \rangle \sigma_n)^\top \\ &= w - \text{ddiag}(w \sigma^\top) \sigma, \end{aligned}$$

where $\text{ddiag} : \mathbb{R}^{n \times n} \rightarrow \mathbb{R}^{n \times n}$ sets all off-diagonal entries of its argument to zero, while leaving diagonal entries the same.

Using this notation and standard tools from matrix manifolds [AMS07], we obtain the Riemannian gradient of f as follows

$$\text{grad} f(\sigma) = P^\perp(\nabla f(\sigma)) = 2(A - \Lambda) \sigma,$$

where $\Lambda = \text{ddiag}(A \sigma \sigma^\top)$. Opening up the terms in the above equality, we obtain

$$\begin{aligned} \text{grad} f(\sigma) &= \left(A - \text{ddiag} \left(A \begin{bmatrix} 1 & \langle \sigma_1, \sigma_2 \rangle & \dots & \langle \sigma_1, \sigma_n \rangle \\ \langle \sigma_2, \sigma_1 \rangle & 1 & \dots & \langle \sigma_2, \sigma_n \rangle \\ \vdots & \vdots & \ddots & \vdots \\ \langle \sigma_n, \sigma_1 \rangle & \langle \sigma_n, \sigma_2 \rangle & \dots & 1 \end{bmatrix} \right) \right) \sigma, \\ &= \left(A - \begin{bmatrix} \langle \sigma_1, g_1 \rangle & & & \\ & \ddots & & \\ & & \langle \sigma_n, g_n \rangle & \\ & & & \end{bmatrix} \right) \sigma \\ &= \begin{bmatrix} -\langle \sigma_1, g_1 \rangle & A_{12} & \dots & A_{1n} \\ A_{21} & -\langle \sigma_2, g_2 \rangle & \dots & A_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ A_{n1} & A_{n2} & \dots & -\langle \sigma_n, g_n \rangle \end{bmatrix} \sigma. \end{aligned}$$

Hence, the Riemannian gradient can be explicitly written as follows

$$\text{grad} f(\sigma) = (g_1 - \langle \sigma_1, g_1 \rangle \sigma_1, \dots, g_n - \langle \sigma_n, g_n \rangle \sigma_n)^\top.$$

In particular, we have

$$\|\text{grad} f(\sigma)\|_{\text{F}}^2 = \sum_{i=1}^n \|g_i - \langle \sigma_i, g_i \rangle \sigma_i\|^2 = \sum_{i=1}^n \left(\|g_i\|^2 - \langle \sigma_i, g_i \rangle^2 \right). \quad (8)$$

Using the same approach, we can calculate the Riemannian Hessian of $f(\sigma)$ along the direction of a vector $u \in T_\sigma \mathcal{M}_r$ by projecting the directional derivative of the gradient vector field onto the tangent space of σ as follows

$$\text{Hess} f(\sigma)[u] = P^\perp(D \text{grad} f(\sigma)[u]),$$

where $D \text{grad} f(\sigma)[u]$ denotes the directional gradient of $\text{grad} f(\sigma)$ along the direction u . This yields

$$\text{Hess} f(\sigma)[u] = P^\perp(2(A - \Lambda)u - 2 \text{ddiag}(A \sigma u^\top + A u \sigma^\top) \sigma) = P^\perp(2(A - \Lambda)u).$$

In particular, we have

$$\langle u, \text{Hess} f(\sigma)[u] \rangle = 2 \langle u, (A - \Lambda)u \rangle, \quad (9)$$

for any $u \in T_\sigma \mathcal{M}_r$.

B Proof of Lemma 3.1

According to the update rule of the BCM algorithm, we have $\sigma_{i_k}^{k+1} = \frac{g_{i_k}^k}{\|g_{i_k}^k\|}$, which yields

$$\begin{aligned}
f(\sigma^{k+1}) &= \sum_{i=1}^n \langle \sigma_i^{k+1}, g_i^{k+1} \rangle \\
&= \langle \sigma_{i_k}^{k+1}, g_{i_k}^{k+1} \rangle + \sum_{i \neq i_k} \langle \sigma_i^{k+1}, g_i^{k+1} \rangle \\
&= \|g_{i_k}^k\| + \sum_{i \neq i_k} \langle \sigma_i^k, g_i^k - A_{ii_k} \sigma_{i_k}^k + A_{ii_k} \frac{g_{i_k}^k}{\|g_{i_k}^k\|} \rangle, \tag{10}
\end{aligned}$$

where the last equality follows since $g_{i_k}^{k+1} = g_{i_k}^k$ as this quantity is independent of $\sigma_{i_k}^{k+1}$ and we have the following for all $i \neq i_k$:

$$\begin{aligned}
g_i^{k+1} &= \sum_{j \neq i} A_{ij} \sigma_j^{k+1} \\
&= A_{ii_k} \sigma_{i_k}^{k+1} + \sum_{j \neq i, i_k} A_{ij} \sigma_j^{k+1} \\
&= A_{ii_k} \frac{g_{i_k}^k}{\|g_{i_k}^k\|} + \sum_{j \neq i, i_k} A_{ij} \sigma_j^k \\
&= A_{ii_k} \frac{g_{i_k}^k}{\|g_{i_k}^k\|} + g_i^k - A_{ii_k} \sigma_{i_k}^k.
\end{aligned}$$

Separating the terms in the sum in (10) and using the fact that A is a symmetric matrix, we obtain

$$\begin{aligned}
f(\sigma^{k+1}) &= \|g_{i_k}^k\| + \sum_{i \neq i_k} A_{ii_k} \langle \sigma_i^k, \frac{g_{i_k}^k}{\|g_{i_k}^k\|} - \sigma_{i_k}^k \rangle + \sum_{i \neq i_k} \langle \sigma_i^k, g_i^k \rangle \\
&= \|g_{i_k}^k\| + \langle g_{i_k}^k, \frac{g_{i_k}^k}{\|g_{i_k}^k\|} - \sigma_{i_k}^k \rangle + \sum_{i \neq i_k} \langle \sigma_i^k, g_i^k \rangle \\
&= 2 \|g_{i_k}^k\| - \langle \sigma_{i_k}^k, g_{i_k}^k \rangle + \sum_{i \neq i_k} \langle \sigma_i^k, g_i^k \rangle \\
&= f(\sigma^k) + 2 (\|g_{i_k}^k\| - \langle \sigma_{i_k}^k, g_{i_k}^k \rangle),
\end{aligned}$$

which concludes the proof of the lemma.

C Proof of Theorem 3.2

From Lemma 3.1, we have

$$\begin{aligned}
f(\sigma^{k+1}) - f(\sigma^k) &= 2 (\|g_{i_k}^k\| - \langle \sigma_{i_k}^k, g_{i_k}^k \rangle) \\
&= \frac{2 \|g_{i_k}^k\| (\|g_{i_k}^k\| - \langle \sigma_{i_k}^k, g_{i_k}^k \rangle)}{\|g_{i_k}^k\|} \\
&\geq \frac{\|g_{i_k}^k\|^2 - \langle \sigma_{i_k}^k, g_{i_k}^k \rangle^2}{\|g_{i_k}^k\|}, \tag{11}
\end{aligned}$$

where the inequality follows since $\|g_{i_k}^k\| \geq \langle \sigma_{i_k}^k, g_{i_k}^k \rangle$, for all $\sigma_{i_k}^k \in \mathbb{R}^{n \times r}$. Letting \mathbb{E}_k denote the expectation over i_k given σ^k , we have

$$\mathbb{E}_k f(\sigma^{k+1}) - f(\sigma^k) \geq \sum_{i=1}^n p_i \frac{\|g_i^k\|^2 - \langle \sigma_i^k, g_i^k \rangle^2}{\|g_{i_k}^k\|}.$$

In particular, when $p_i = \frac{1}{n}$, for all $i \in [n]$, we have

$$\mathbb{E}_k f(\sigma^{k+1}) - f(\sigma^k) \geq \frac{1}{n \|A\|_1} \sum_{i=1}^n \left(\|g_i^k\|^2 - \langle \sigma_i^k, g_i^k \rangle^2 \right),$$

since $\|g_{i_k}^k\|$ can be upper bounded as follows

$$\|g_{i_k}^k\| \leq \sum_{j \neq i_k} |A_{i_k j}| \|\sigma_j^k\| \leq \|A\|_1.$$

Therefore, we have

$$\mathbb{E}_k f(\sigma^{k+1}) - f(\sigma^k) \geq \frac{\|\text{grad} f(\sigma^k)\|_F^2}{n \|A\|_1}.$$

On the other hand, when $p_i = \frac{\|g_i^k\|}{\sum_{j=1}^n \|g_j^k\|}$, we have

$$\mathbb{E}_k f(\sigma^{k+1}) - f(\sigma^k) \geq \frac{1}{\|A\|_{1,1}} \sum_{i=1}^n \left(\|g_i^k\|^2 - \langle \sigma_i^k, g_i^k \rangle^2 \right),$$

where $\|A\|_{1,1} = \sum_{i,j=1}^n |A_{ij}|$ is the $L_{1,1}$ norm of matrix A . This yields

$$\mathbb{E}_k f(\sigma^{k+1}) - f(\sigma^k) \geq \frac{\|\text{grad} f(\sigma^k)\|_F^2}{\|A\|_{1,1}}. \quad (12)$$

For simplicity, in the rest of the proof we consider the latter case, yet replacing $\|A\|_{1,1}$ by $n \|A\|_1$ in what follows, we get the convergence rate guarantee for uniform probability case as well.

In order to prove (2), we assume the contrary that $\mathbb{E} \|\text{grad} f(\sigma^k)\|_F^2 > \epsilon$, for all $k \in [K-1]$. Then, using the boundedness of f , we get

$$f^* - f(\sigma^0) \geq \mathbb{E} f(\sigma^K) - f(\sigma^0) = \sum_{k=0}^{K-1} \mathbb{E} [f(\sigma^{k+1}) - f(\sigma^k)] = \sum_{k=0}^{K-1} \mathbb{E} [\mathbb{E}_k f(\sigma^{k+1}) - f(\sigma^k)].$$

Using the expected functional ascent of BCM in (12) above, we get

$$f^* - f(\sigma^0) \geq \sum_{k=0}^{K-1} \frac{\mathbb{E} \|\text{grad} f(\sigma^k)\|_F^2}{\|A\|_{1,1}} > \frac{K\epsilon}{\|A\|_{1,1}},$$

where the last inequality follows by the assumption. Then, by contradiction, the algorithm returns a solution with $\mathbb{E} \|\text{grad} f(\sigma^k)\|_F^2 \leq \epsilon$, for some $k \in [K-1]$, provided that

$$K \geq \frac{\|A\|_{1,1} (f^* - f(\sigma^0))}{\epsilon}.$$

D Proof of Lemma 4.1

By Lemma 3.1, we have

$$f(\sigma^{k+1}) - f(\sigma^k) = 2 \left(\|g_{i_k}^k\| - \langle \sigma_{i_k}^k, g_{i_k}^k \rangle \right) \geq 0.$$

Since $f(\sigma) \leq \|A\|_{1,1} < \infty$, for any $\sigma \in \mathcal{M}_r$, then $\{f(\sigma^k)\}_{k \geq 0}$ converges. Hence, letting $\bar{\sigma}$ be a limit point of the sequence $\{\sigma^k\}_{k \geq 0}$, we can conclude that $\bar{\sigma} \in \mathcal{M}_r$ since \mathcal{M}_r is a closed manifold. Since any fixed point σ of the BCM algorithm satisfies $\sigma_i = \frac{g_i}{\|g_i\|}$ and the limit points of the sequence $\{\sigma^k\}_{k \geq 0}$ are fixed points of the BCM algorithm, the lemma follows.

E Proof of Theorem 4.2

By (12), we have

$$\mathbb{E}_k f(\sigma^{k+1}) - f(\sigma^k) \geq \frac{\|\text{grad}f(\sigma^k)\|_{\mathbb{F}}^2}{\|A\|_{1,1}}. \quad (13)$$

Our aim is to show that $\|\text{grad}f(\sigma^k)\|_{\mathbb{F}}^2 \geq c(\bar{f} - f(\sigma^k))$, for some $0 < c < \|A\|_{1,1}$, in a neighborhood around the limit points of the iterates generated by the algorithm. To this end, we consider the following formulation

$$\sigma_i^k = \sigma_i \cos(\|u_i\| t) + \frac{u_i}{\|u_i\|} \sin(\|u_i\| t), \quad (14)$$

for some $t \in \mathbb{R}$ and $\sigma \in \mathcal{M}_r$ such that $u \in \mathcal{H}_\sigma \bar{\mathcal{M}}_r$ with $\|u\|_{\mathbb{F}} = 1$ (without loss of generality) is the unique horizontal lift (although the particular value of $\sigma \in \mathcal{M}_r$ is not important for our proof, it can be found as the projection of σ^k onto the set $\{\bar{\sigma}Q : Q \in O(r)\}$). The second order Taylor approximation to (14) yields

$$\sigma_i^k = \sigma_i + tu_i - \frac{t^2}{2} \|u_i\|^2 \sigma_i + O(t^3).$$

Using this approximation, we obtain

$$g_i^k = g_i + tv_i - \frac{t^2}{2} \tilde{g}_i + O(t^3),$$

where

$$v_i^k = \sum_{j \neq i} A_{ij} u_j \quad \text{and} \quad \tilde{g}_i = \sum_{j \neq i} A_{ij} \|u_j\|^2 \sigma_j.$$

This yields the following Taylor approximation to $\|\text{grad}f(\sigma^k)\|_{\mathbb{F}}^2$:

$$\begin{aligned} \|\text{grad}f(\sigma^k)\|_{\mathbb{F}}^2 &= \sum_{i=1}^n \left(\|g_i^k\|^2 - \langle \sigma_i^k, g_i^k \rangle^2 \right) \\ &= \sum_{i=1}^n \left(\left\| g_i + tv_i - \frac{t^2}{2} \tilde{g}_i \right\|^2 - \left\langle \sigma_i + tu_i - \frac{t^2}{2} \|u_i\|^2 \sigma_i, g_i + tv_i - \frac{t^2}{2} \tilde{g}_i \right\rangle^2 \right) + O(t^3) \\ &= \sum_{i=1}^n \left\{ \|g_i\|^2 + 2t \langle g_i, v_i \rangle - t^2 \langle g_i, \tilde{g}_i \rangle + t^2 \|v_i\|^2 \right. \\ &\quad \left. - \left(\langle \sigma_i, g_i \rangle + t \langle \sigma_i, v_i \rangle - \frac{t^2}{2} \langle \sigma_i, \tilde{g}_i \rangle + t \langle u_i, g_i \rangle + t^2 \langle u_i, v_i \rangle - \frac{t^2}{2} \|u_i\|^2 \langle \sigma_i, g_i \rangle \right)^2 \right\} + O(t^3). \end{aligned}$$

Note that $\bar{\sigma}$ is a limit point of the algorithm that satisfy $\bar{\sigma}_i = \frac{\tilde{g}_i}{\|\tilde{g}_i\|}$. Since $\bar{\sigma} = \sigma Q$, for some rotation matrix $Q \in O(r)$, then we have $\sigma_i = \frac{g_i}{\|g_i\|}$. Since $u \in T_\sigma \mathcal{M}_r$, we also have $\langle \sigma_i, u_i \rangle = 0$. Using these relations in the above equality yields

$$\begin{aligned} \|\text{grad}f(\sigma^k)\|_{\mathbb{F}}^2 &= \sum_{i=1}^n \left\{ \|g_i\|^2 + 2t \|g_i\| \langle \sigma_i, v_i \rangle - t^2 \|g_i\| \langle \sigma_i, \tilde{g}_i \rangle + t^2 \|v_i\|^2 \right. \\ &\quad \left. - \left(\|g_i\| + t \langle \sigma_i, v_i \rangle - \frac{t^2}{2} \langle \sigma_i, \tilde{g}_i \rangle + t^2 \langle u_i, v_i \rangle - \frac{t^2}{2} \|u_i\|^2 \|g_i\| \right)^2 \right\} + O(t^3) \\ &= \sum_{i=1}^n \left\{ \|g_i\|^2 + 2t \|g_i\| \langle \sigma_i, v_i \rangle - t^2 \|g_i\| \langle \sigma_i, \tilde{g}_i \rangle + t^2 \|v_i\|^2 \right. \\ &\quad \left. - \left(\|g_i\|^2 + 2t \|g_i\| \langle \sigma_i, v_i \rangle - t^2 \|g_i\| \langle \sigma_i, \tilde{g}_i \rangle + 2t^2 \|g_i\| \langle u_i, v_i \rangle \right. \right. \\ &\quad \left. \left. - t^2 \|u_i\|^2 \|g_i\|^2 + t^2 \langle \sigma_i, v_i \rangle^2 \right) \right\} + O(t^3) \\ &= t^2 \sum_{i=1}^n \left(\|v_i\|^2 - \langle \sigma_i, v_i \rangle^2 - 2 \|g_i\| \langle u_i, v_i \rangle + \|u_i\|^2 \|g_i\|^2 \right) + O(t^3). \quad (15) \end{aligned}$$

Since $\langle \sigma_i, u_i \rangle = 0$, we have by the Pythagorean theorem that

$$\|v_i\|^2 - \langle \sigma_i, v_i \rangle^2 - \left\langle \frac{u_i}{\|u_i\|}, v_i \right\rangle^2 \geq 0.$$

Using this inequality in (15), we get

$$\begin{aligned} \|\text{grad} f(\sigma^k)\|_{\mathbb{F}}^2 &\geq t^2 \sum_{i=1}^n \left(\left\langle \frac{u_i}{\|u_i\|}, v_i \right\rangle^2 - 2 \|g_i\| \langle u_i, v_i \rangle + \|u_i\|^2 \|g_i\|^2 \right) + O(t^3) \\ &= t^2 \sum_{i=1}^n \left(\|u_i\| \|g_i\| - \left\langle \frac{u_i}{\|u_i\|}, v_i \right\rangle \right)^2 + O(t^3). \end{aligned} \quad (16)$$

In order to lower bound (16) by $c(f(\sigma) - f(\sigma^k))$, we consider the second order Taylor approximation of $f(\sigma^k)$, which can be written as follows

$$\begin{aligned} f(\sigma^k) &= \sum_{i=1}^n \langle \sigma_i^k, g_i^k \rangle \\ &= \sum_{i=1}^n \left\langle \sigma_i + tu_i - \frac{t^2}{2} \|u_i\|^2 \sigma_i, g_i + tv_i - \frac{t^2}{2} \tilde{g}_i \right\rangle + O(t^3) \\ &= \sum_{i=1}^n \left(\langle \sigma_i, g_i \rangle + t \langle \sigma_i, v_i \rangle - \frac{t^2}{2} \langle \sigma_i, \tilde{g}_i \rangle + t \langle u_i, g_i \rangle + t^2 \langle u_i, v_i \rangle - \frac{t^2}{2} \|u_i\|^2 \langle \sigma_i, g_i \rangle \right) + O(t^3). \end{aligned}$$

Similar to the previous derivations, using the fact that $\sigma_i = \frac{g_i}{\|g_i\|}$ and $\langle \sigma_i, u_i \rangle = 0$, for all $i \in [n]$, we obtain

$$\begin{aligned} f(\sigma^k) &= f(\sigma) + \sum_{i=1}^n \left(t \langle \sigma_i, v_i \rangle - \frac{t^2}{2} \langle \sigma_i, \tilde{g}_i \rangle + t^2 \langle u_i, v_i \rangle - \frac{t^2}{2} \|u_i\|^2 \langle \sigma_i, g_i \rangle \right) + O(t^3) \\ &= f(\sigma) + \sum_{i=1}^n \left(t \sum_{j \neq i} A_{ij} \langle \sigma_i, u_j \rangle - \frac{t^2}{2} \sum_{j \neq i} A_{ij} \|u_j\|^2 \langle \sigma_i, \sigma_j \rangle + t^2 \langle u_i, v_i \rangle - \frac{t^2}{2} \|u_i\|^2 \langle \sigma_i, g_i \rangle \right) + O(t^3) \\ &= f(\sigma) + t \sum_{j=1}^n \sum_{i \neq j} A_{ji} \langle \sigma_i, u_j \rangle - \frac{t^2}{2} \sum_{j=1}^n \sum_{i \neq j} A_{ji} \|u_j\|^2 \langle \sigma_i, \sigma_j \rangle \\ &\quad + t^2 \sum_{i=1}^n \left(\langle u_i, v_i \rangle - \frac{1}{2} \|u_i\|^2 \langle \sigma_i, g_i \rangle \right) + O(t^3), \end{aligned}$$

where the last line follows since A is symmetric. Using the definition $g_j = \sum_{i \neq j} A_{ji} \sigma_i$ and $\sigma_i = \frac{g_i}{\|g_i\|}$ in the above inequality yields

$$\begin{aligned} f(\sigma^k) &= f(\sigma) + t \sum_{j=1}^n \langle g_j, u_j \rangle - \frac{t^2}{2} \sum_{j=1}^n \|u_j\|^2 \langle g_j, \sigma_j \rangle + t^2 \sum_{i=1}^n \left(\langle u_i, v_i \rangle - \frac{1}{2} \|u_i\|^2 \langle \sigma_i, g_i \rangle \right) + O(t^3) \\ &= f(\sigma) + t^2 \sum_{i=1}^n \left(\langle u_i, v_i \rangle - \|u_i\|^2 \|g_i\| \right) + O(t^3). \end{aligned}$$

Reorganizing terms, we get

$$\bar{f} - f(\sigma^k) = f(\sigma) - f(\sigma^k) = t^2 \sum_{i=1}^n \left(\|u_i\|^2 \|g_i\| - \langle u_i, v_i \rangle \right) + O(t^3). \quad (17)$$

Turning back our attention to (16), we can lower bound the right-hand side as follows

$$\begin{aligned}
\|\text{grad}f(\sigma^k)\|_{\mathbb{F}}^2 &\geq t^2 \sum_{i=1}^n \frac{1}{\|u_i\|^2} \left(\|u_i\|^2 \|g_i\| - \langle u_i, v_i \rangle \right)^2 + O(t^3) \\
&\geq t^2 \sum_{i=1}^n \left(\|u_i\|^2 \|g_i\| - \langle u_i, v_i \rangle \right)^2 + O(t^3) \\
&\geq \frac{t^2}{n} \left(\sum_{i=1}^n \left(\|u_i\|^2 \|g_i\| - \langle u_i, v_i \rangle \right) \right)^2 + O(t^3),
\end{aligned}$$

where the second inequality follows since $\|u_i\|^2 \leq \|u\|_{\mathbb{F}}^2 = 1$ and the last inequality follows since $(\sum_{i=1}^n a_i)^2 \leq n \sum_{i=1}^n a_i^2$, for all $a_i \in \mathbb{R}$, $i \in [n]$. Using the second order approximation derived in (17) in the above inequality, we obtain

$$\begin{aligned}
\|\text{grad}f(\sigma^k)\|_{\mathbb{F}}^2 &\geq \frac{\bar{f} - f(\sigma^k)}{n} \sum_{i=1}^n \left(\|u_i\|^2 \|g_i\| - \langle u_i, v_i \rangle \right) + O(t^3) \\
&= \frac{\langle u, (\Lambda - A)u \rangle}{n} (\bar{f} - f(\sigma^k)) + O(t^3),
\end{aligned}$$

where $\Lambda = \text{diag}(\|g_1\|, \dots, \|g_n\|)$. Since we have $\langle u, (A - \Lambda)u \rangle \leq -\mu \|u\|_{\mathbb{F}}^2$, for all $u \in T_{\sigma} \mathcal{M}_r$ and $\langle u, \pi^{(\ell)} \rangle$, for all $\ell \in [k-1]$, we conclude that

$$\|\text{grad}f(\sigma^k)\|_{\mathbb{F}}^2 \geq \frac{\mu}{n} (\bar{f} - f(\sigma^k)) + O(t^3).$$

Since the above Taylor approximation is made around the set of limit points of the sequence $\{\sigma^k\}_{k \geq 0}$, there exists an integer $K > 0$ such that for all $k \geq K$, t is small enough to satisfy $O(t^3) \geq -\delta_K (\bar{f} - f(\sigma^k))$ in the above inequality, for some $\delta_K < \frac{\mu}{n}$. Therefore, we have

$$\|\text{grad}f(\sigma^k)\|_{\mathbb{F}}^2 \geq \left(\frac{\mu}{n} - \delta_K \right) (\bar{f} - f(\sigma^k)). \quad (18)$$

for all $k \geq K$. Combining this inequality with (13), we get

$$\mathbb{E}_k f(\sigma^{k+1}) - f(\sigma^k) \geq \left(\frac{\mu}{n \|A\|_{1,1}} - \delta_K \right) (\bar{f} - f(\sigma^k)),$$

for all $k \geq K$, with a slight abuse of notation in the definition of the term δ_K . Rearranging terms in the above inequality concludes the proof.