

Learning with Maximum Likelihood



René Magritte, "La reproduction interdite" (1937)

Likelihood: Bernoulli Variables

- Suppose a fair coin is tossed n times, independently
 - $Y \sim \text{Bernoulli}(\theta)$
- The likelihood (discrete case) is the probability of observing the dataset when the parameters are θ
 - $P(Y_i = 1|\theta) = \theta$
 - $P(Y_i = 0|\theta) = 1 - \theta$
 - $P(Y_i = y_i|\theta) = \theta^{y_i}(1 - \theta)^{1-y_i}$
- $P(Y_1 = y_1, Y_2 = y_2, \dots, Y_m = y_m|\theta) = \prod_{i=1}^m P(Y_i = y_i|\theta)$

Maximum likelihood: Bernoulli

- Suppose we observe the data $Y^{(1)} = y^{(1)}, Y^{(2)} = y^{(2)}, \dots, Y^{(m)} = y^{(m)}$ (m i.i.d. Bernoulli variables), and would like to know what θ is
- One possibility: find the θ that maximizes the likelihood function
 - What value of θ makes the data set that we are actually observing (i.e., the training set) the most plausible?
- $P(Y_1 = y_1, Y_2 = y_2, \dots, Y_m = y_m | \theta)$ is maximized at $\theta = \frac{1}{m} \sum_{i=1}^m y_i$

Likelihood: Gaussian Noise

- Assume each data point is generated using some process.

- E.g., $y^{(i)} = \theta^T x^{(i)} + \epsilon^{(i)}$, $\epsilon^{(i)} \sim N(0, \sigma^2)$

- We can now compute the likelihood of single datapoint

- I.e., the probability of the point for a set θ .

- E.g., $P(y^{(i)} | \theta, x^{(i)}) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(y^{(i)} - \theta^T x^{(i)})^2}{2\sigma^2}\right)$ We can then compute the likelihood for the entire training set $\{(x^{(1)}, y^{(1)}), (x^{(2)}, y^{(2)}), \dots, (x^{(m)}, y^{(m)})\}$ (assuming each point is independent)

- E.g., $P(y | \theta, x) = \prod_{i=1}^m \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(y^{(i)} - \theta^T x^{(i)})^2}{2\sigma^2}\right)$

Maximum Likelihood

- $P(\text{data}|\theta) = P(y|\theta, x) = \prod_1^m \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(y^{(i)} - \theta^T x^{(i)})^2}{2\sigma^2}\right)$
- $\log P(\text{data}|\theta) = \sum -\frac{(y^{(i)} - \theta^T x^{(i)})^2}{2\sigma^2} + 2m/\log(2\pi\sigma^2)$

is maximized for a value of θ for which $\sum_{i=1}^m (y^{(i)} - \theta^T x^{(i)})^2$ is minimized
- Note: x is fixed