

# Linear Algebra - Part II

## Projection, Eigendecomposition, SVD

Punit Shah

(Adapted from Sargur Srihari's [slides](#))

## Brief Review from Part 1

- ▶ Symmetric Matrix:

$$\mathbf{A} = \mathbf{A}^T$$

- ▶ Orthogonal Matrix:

$$\mathbf{A}^T \mathbf{A} = \mathbf{A} \mathbf{A}^T = \mathbf{I} \quad \text{and} \quad \mathbf{A}^{-1} = \mathbf{A}^T$$


- ▶ L2 Norm:

$$\|\mathbf{x}\|_2 = \sqrt{\sum_i x_i^2}$$

## Angle Between Vectors

- ▶ Dot product of two vectors can be written in terms of their L2 norms and the angle  $\theta$  between them.

$$\mathbf{a}^T \mathbf{b} = \|\mathbf{a}\|_2 \|\mathbf{b}\|_2 \cos(\theta)$$



# Cosine Similarity

- ▶ Cosine between two vectors is a measure of their similarity:

$$\cos(\theta) = \frac{\mathbf{a} \cdot \mathbf{b}}{\|\mathbf{a}\| \|\mathbf{b}\|}$$

- ▶ **Orthogonal Vectors:** Two vectors  $\mathbf{a}$  and  $\mathbf{b}$  are orthogonal to each other if  $\mathbf{a} \cdot \mathbf{b} = 0$ .

## Vector Projection

- ▶ Given two vectors  $\mathbf{a}$  and  $\mathbf{b}$ , let  $\hat{\mathbf{b}} = \frac{\mathbf{b}}{\|\mathbf{b}\|}$  be the unit vector in the direction of  $\mathbf{b}$ .
- ▶ Then  $\mathbf{a}_1 = a_1 \hat{\mathbf{b}}$  is the orthogonal projection of  $\mathbf{a}$  onto a straight line parallel to  $\mathbf{b}$ , where

$$a_1 = \|\mathbf{a}\| \cos(\theta) = \mathbf{a} \cdot \hat{\mathbf{b}} = \mathbf{a} \cdot \frac{\mathbf{b}}{\|\mathbf{b}\|}$$




Image taken from [wikipedia](#).

## Diagonal Matrix

- ▶ Diagonal matrix has mostly zeros with non-zero entries only in the diagonal, e.g. identity matrix.
- ▶ A square diagonal matrix with diagonal elements given by entries of vector  $\mathbf{v}$  is denoted:

$$\text{diag}(\mathbf{v})$$

- ▶ Multiplying vector  $\mathbf{x}$  by a diagonal matrix is efficient:

$$\text{diag}(\mathbf{v})\mathbf{x} = \mathbf{v} \odot \mathbf{x}$$

- ▶ Inverting a square diagonal matrix is efficient:

$$\text{diag}(\mathbf{v})^{-1} = \text{diag}\left(\left[\frac{1}{v_1}, \dots, \frac{1}{v_n}\right]^T\right)$$

# Determinant

- ▶ Determinant of a square matrix is a mapping to a scalar.

$$\det(\mathbf{A}) \quad \text{or} \quad |\mathbf{A}|$$

- ▶ Measures how much multiplication by the matrix expands or contracts the space.
- ▶ Determinant of product is the product of determinants:

$$\det(\mathbf{AB}) = \det(\mathbf{A})\det(\mathbf{B})$$

$$|A| = \begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc.$$

*Image taken from [wikipedia](#).*

# List of Equivalencies

The following are all equivalent:

- ▶  $\mathbf{A}$  is invertible, i.e.  $\mathbf{A}^{-1}$  exists.
- ▶  $\mathbf{Ax} = \mathbf{b}$  has a **unique** solution.
- ▶ Columns of  $\mathbf{A}$  are linearly independent.
- ▶  $\det(\mathbf{A}) \neq 0$
- ▶  $\mathbf{Ax} = \mathbf{0}$  has a unique, trivial solution:  $\mathbf{x} = \mathbf{0}$ .

# Zero Determinant

If  $\det(\mathbf{A}) = 0$ , then:

- ▶  $\mathbf{A}$  is linearly dependent.
- ▶  $\mathbf{Ax} = \mathbf{b}$  has no solution or infinitely many solutions.
- ▶  $\mathbf{Ax} = \mathbf{0}$  has a non-zero solution.

# Matrix Decomposition

- ▶ We can decompose an integer into its prime factors, e.g.  
 $12 = 2 \times 2 \times 3$ .
- ▶ Similarly, matrices can be decomposed into factors to learn universal properties:

$$\mathbf{A} = \mathbf{V}\text{diag}(\boldsymbol{\lambda})\mathbf{V}^{-1}$$

# Eigenvectors

- ▶ An eigenvector of a square matrix  $\mathbf{A}$  is a nonzero vector  $\mathbf{v}$  such that multiplication by  $\mathbf{A}$  only changes the scale of  $\mathbf{v}$ .

$$\mathbf{A}\mathbf{v} = \lambda\mathbf{v}$$

- ▶ The scalar  $\lambda$  is known as the **eigenvalue**.
- ▶ If  $\mathbf{v}$  is an eigenvector of  $\mathbf{A}$ , so is any rescaled vector  $s\mathbf{v}$ . Moreover,  $s\mathbf{v}$  still has the same eigenvalue. Thus, we constrain the eigenvector to be of unit length:

$$\|\mathbf{v}\| = 1$$

# Characteristic Polynomial

- ▶ Eigenvalue equation of matrix  $\mathbf{A}$ :

$$\mathbf{A}\mathbf{v} = \lambda\mathbf{v}$$

$$\mathbf{A}\mathbf{v} - \lambda\mathbf{v} = \mathbf{0}$$

$$(\mathbf{A} - \lambda\mathbf{I})\mathbf{v} = \mathbf{0}$$

- ▶ If nonzero solution for  $\mathbf{v}$  exists, then it must be the case that:

$$\det(\mathbf{A} - \lambda\mathbf{I}) = 0$$

- ▶ Unpacking the determinant as a function of  $\lambda$ , we get:

$$|\mathbf{A} - \lambda\mathbf{I}| = (\lambda_1 - \lambda)(\lambda_2 - \lambda)\dots(\lambda_n - \lambda) = 0$$

- ▶ The  $\lambda_1, \lambda_2, \dots, \lambda_n$  are roots of the characteristic polynomial, and are eigenvalues of  $\mathbf{A}$ .

## Example

- ▶ Consider the matrix:

$$\mathbf{A} = \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix}$$

- ▶ The characteristic polynomial is:

$$\det(\mathbf{A} - \lambda\mathbf{I}) = \det \begin{bmatrix} 2 - \lambda & 1 \\ 1 & 2 - \lambda \end{bmatrix} = 3 - 4\lambda + \lambda^2 = 0$$

- ▶ It has roots  $\lambda = 1$  and  $\lambda = 3$  which are the two eigenvalues of  $\mathbf{A}$ .
- ▶ We can then solve for eigenvectors using  $\mathbf{A}\mathbf{v} = \lambda\mathbf{v}$ :

$$\mathbf{v}_{\lambda=1} = [1, -1]^T \quad \text{and} \quad \mathbf{v}_{\lambda=3} = [1, 1]^T$$

# Eigendecomposition

- ▶ Suppose that  $n \times n$  matrix  $\mathbf{A}$  has  $n$  linearly independent eigenvectors  $\{\mathbf{v}^{(1)}, \dots, \mathbf{v}^{(n)}\}$  with eigenvalues  $\{\lambda_1, \dots, \lambda_n\}$ .
- ▶ Concatenate eigenvectors to form matrix  $\mathbf{V}$ .
- ▶ Concatenate eigenvalues to form vector  $\boldsymbol{\lambda} = [\lambda_1, \dots, \lambda_n]^T$ .
- ▶ The **eigendecomposition** of  $\mathbf{A}$  is given by:

$$\mathbf{A} = \mathbf{V}\text{diag}(\boldsymbol{\lambda})\mathbf{V}^{-1}$$


# Symmetric Matrices

- ▶ Every real symmetric matrix  $\mathbf{A}$  can be decomposed into real-valued eigenvectors and eigenvalues:

$$\mathbf{A} = \mathbf{Q}\mathbf{\Lambda}\mathbf{Q}^T$$


- ▶  $\mathbf{Q}$  is an orthogonal matrix of the eigenvectors of  $\mathbf{A}$ , and  $\mathbf{\Lambda}$  is a diagonal matrix of eigenvalues.
- ▶ We can think of  $\mathbf{A}$  as scaling space by  $\lambda_i$  in direction  $\mathbf{v}^{(i)}$ .

Plot of unit vectors  $\mathbf{u} \in \mathbb{R}^2$   
(circle)



with two variables  $x_1$  and  $x_2$

Plot of vectors  $\mathbf{A}\mathbf{u}$   
(ellipse)



## Eigendecomposition is not Unique

- ▶ Decomposition is not unique when two eigenvalues are the same.
- ▶ By convention, order entries of  $\Lambda$  in descending order. Then, eigendecomposition is unique if all eigenvalues are unique.
- ▶ If any eigenvalue is zero, then the matrix is **singular**.

# Positive Definite Matrix

- ▶ A matrix whose eigenvalues are all positive is called **positive definite**.
- ▶ If eigenvalues are positive or zero, then matrix is called **positive semidefinite**.
- ▶ Positive definite matrices guarantee that:

$$\mathbf{x}^T \mathbf{A} \mathbf{x} > 0 \quad \text{for any nonzero vector } \mathbf{x}$$

- ▶ Similarly, positive semidefinite guarantees:  $\mathbf{x}^T \mathbf{A} \mathbf{x} \geq 0$

# Singular Value Decomposition (SVD)

- ▶ If  $\mathbf{A}$  is not square, eigendecomposition is undefined.
- ▶ SVD is a decomposition of the form:

$$\mathbf{A} = \mathbf{U}\mathbf{D}\mathbf{V}^T$$

- ▶ SVD is more general than eigendecomposition.
- ▶ Every real matrix has a SVD.

## SVD Definition (1)

- ▶ Write  $\mathbf{A}$  as a product of three matrices:  $\mathbf{A} = \mathbf{U}\mathbf{D}\mathbf{V}^T$ .
- ▶ If  $\mathbf{A}$  is  $m \times n$ , then  $\mathbf{U}$  is  $m \times m$ ,  $\mathbf{D}$  is  $m \times n$ , and  $\mathbf{V}$  is  $n \times n$ .
- ▶  $\mathbf{U}$  and  $\mathbf{V}$  are orthogonal matrices, and  $\mathbf{D}$  is a diagonal matrix (not necessarily square).
- ▶ Diagonal entries of  $\mathbf{D}$  are called **singular values** of  $\mathbf{A}$ .
- ▶ Columns of  $\mathbf{U}$  are the **left singular vectors**, and columns of  $\mathbf{V}$  are the **right singular vectors**.

## SVD Definition (2)

---

- ▶ SVD can be interpreted in terms of eigendecomposition.
- ▶ Left singular vectors of  $\mathbf{A}$  are the eigenvectors of  $\mathbf{A}\mathbf{A}^T$ .
- ▶ Right singular vectors of  $\mathbf{A}$  are the eigenvectors of  $\mathbf{A}^T\mathbf{A}$ .
- ▶ Nonzero singular values of  $\mathbf{A}$  are square roots of eigenvalues of  $\mathbf{A}^T\mathbf{A}$  and  $\mathbf{A}\mathbf{A}^T$ .