

Digital Photography I

optics and sensors



CSC2529

David Lindell

University of Toronto

cs.toronto.edu/~lindell/teaching/2529

*slides adapted from Gordon Wetzstein,
Fredo Durand, Ioannis Gkioulekas, Marc Levoy

Announcements

- HW 1 is due Friday at 11:59pm
- HW 2 is out (due next Friday 9/27)
- TA office hours Wed 1-2pm BA 5256

Colloquium in Applied AI

featuring speakers from the Technion, Tel Aviv University, the Weizmann Institute and the University of Toronto

Thursday, September 12, 2024

9 a.m. – 5 p.m.*

University of Toronto

Presented by the University of Toronto Department of Computer Science,
in collaboration with the Data Sciences Institute



Michal Irani



Title: Reading Minds & Machines

1. Can we reconstruct images that a person saw, directly from his/her fMRI brain recordings?
2. Can we reconstruct the training data that a deep-network trained on, directly from the parameters of the network?

The answer to both of these intriguing questions is "Yes!"



Toronto Vision Seminar



Scan to register and join the meeting

Michal Irani

Professor of Computer Science and Applied Mathematics

“Deep Internal Learning” – Deep Learning
Without Prior Examples



Friday, Sep 13th



12:00pm–1:00pm



BA3200

Reception to follow

Let's say we have a sensor...



digital sensor
(CCD or
CMOS)

... and an object we like to photograph

real-world
object



digital sensor
(CCD or
CMOS)



What would an image taken like this look like?

Bare-sensor imaging


real-world
object




digital sensor
(CCD or
CMOS)




Bare-sensor imaging



Bare-sensor imaging



Bare-sensor imaging



Bare-sensor imaging



All scene points contribute to all sensor pixels

What can we do to make our image look better?


real-world
object



digital sensor
(CCD or
CMOS)




Let's add something to this scene




What would an image taken like this look like?


Pinhole imaging



Pinhole imaging




Pinhole imaging




Each scene point contributes to only one sensor pixel

What does the image on the sensor look like?


Pinhole imaging



Pinhole camera terms




Pinhole camera terms



Focal length


real-world
object



Focal length


What happens as we change the focal length?

real-world
object




Focal length

What happens as we change the focal length?



Focal length

What happens as we change the focal length?



Pinhole size

real-world
object



pinhole
diameter



Ideal pinhole has infinitesimally small size

- In practice that is impossible.

Pinhole size

What happens as we change the pinhole diameter?

real-world
object




pinhole
diameter



Pinhole size


What happens as we change the pinhole diameter?



Pinhole size

What happens as we change the pinhole diameter?

real-world
object





Pinhole size

What happens as we change the pinhole diameter?


object projection becomes blurrier

real-world
object



Pinhole size


What happens as we change the pinhole diameter?




Will the image keep getting sharper the smaller we make the pinhole?

Diffraction limit

A consequence of the wave nature of light




What do geometric optics
predict will happen?




What do wave optics
predict will happen?

Diffraction limit

A consequence of the wave nature of light




What do geometric optics
predict will happen?




What do wave optics
predict will happen?

Diffraction limit

A consequence of the wave nature of light



What do geometric optics
predict will happen?




What do wave optics
predict will happen?

Diffraction limit


Diffraction pattern = Fourier transform of the pinhole.

- Smaller pinhole means bigger Fourier spectrum.
- Smaller pinhole means more diffraction.



small pinhole

wide
diffraction
pattern



large pinhole


narrow
diffraction
pattern

What about light efficiency?

real-world
object



pinhole
diameter



focal length f


- What is the effect of doubling the pinhole diameter?
- What is the effect of doubling the focal length?

What about light efficiency?

real-world
object




pinhole
diameter





- 2x pinhole diameter \rightarrow 4x light
- 2x focal length \rightarrow $\frac{1}{4}$ x light

Pinhole Camera / Camera Obscura



Mo-Ti (Chinese Philosopher) 470-390 BC






Bryan Bowen
@bryanmbowen



Fun discovery - a small crack in the eastern facade of the Canada Malting Co silos has created a perfect pinhole camera. The result: real time projection of Toronto's waterfront on the silo's interior curved surfaces. An unplugged projection show!



9:37 AM · Jan 27, 2022 · Twitter for iPhone

656 Retweets 70 Quote Tweets 2,836 Likes











Abelardo Morell

Pinhole Camera / Camera Obscura



J. Vermeer "The Milkmaid", 1658



同サイズの《牛乳を注ぐ女》が投影される額縁の前に立ち

Credit: ©Toppan Printing Co., Ltd.

Original photo data (Het melkmeisje [The Milkmaid] by Johannes Vermeer) :


©Rijksmuseum Amsterdam. Purchased with the support of the Vereniging Rembrandt

Digital Photography - Overview


- optics
- aperture
- depth of field
- field of view
- exposure
- noise
- color filter arrays
- image processing pipeline



SLR Camera



SLR Camera



Camera Optics

Niepce "View from the Window at Le Gras", 1826



1826
8h exp

Daguerrotype



- invented in 1836 by Louis Daguerre
- lenses focus light, better chemicals!

Daguerre "Boulevard du Temple", 1838



exposure
10-12 mins

Lenses


- focus light
- magnify objects

Nimrud lens - 2700 years old




What is a lens?

A piece of glass manufactured to have a specific shape




What is a lens?

A piece of glass manufactured to have a specific shape





How does a lens work?




Refraction

Refraction is the bending of rays of light when they move from one material to another



How does a lens work?

Lenses are designed so that their refraction makes light rays bend in a very specific way.




Thin lens model

Simplification of geometric optics for well-designed lenses.



Thin lens model

Simplification of geometric optics for well-designed lenses.




Two assumptions:

1. Rays passing through lens center are unaffected.

Thin lens model

Simplification of geometric optics for well-designed lenses.




Two assumptions:

1. Rays passing through lens center are unaffected.
2. Parallel rays converge to a single point located on focal plane.

Thin lens model

Simplification of geometric optics for well-designed lenses.




Two assumptions:

1. Rays passing through lens center are unaffected.
2. Parallel rays converge to a single point located on focal plane.

Thin lens model

Simplification of geometric optics for well-designed lenses.




Two assumptions:

1. Rays passing through lens center are unaffected.
2. Parallel rays converge to a single point located on focal plane.

Thin Lens Model

Ray tracing example


- Parallel rays map to the focal plane



Thin Lens Model

Ray tracing example


- Parallel rays map to the focal plane
- The chief ray passes straight through the center



Thin Lens Model

Ray tracing example

- Parallel rays map to the focal plane
- The chief ray passes straight through the center
- The ray that passes through the near focal plane becomes parallel




Thin Lens Model

Thin lens formula

$$\frac{1}{f} = \frac{1}{S_1} + \frac{1}{S_2}$$


magnification:

$$M = -\frac{S_2}{S_1} = \frac{f}{f - S_1}$$




Lenses

$S_1 < f$: magnifying glass




Lenses

$S_1 < f$: magnifying glass




minification



Yes, but...

Thin lenses are a fiction


The thin lens model assumes that the lens has no thickness, but this is rarely true...



To make real lenses behave like ideal thin lenses, we have to use combinations of multiple lens elements (compound lenses).

Thin lenses are a fiction





The thin lens model assumes that the lens has no thickness, but this is rarely true...







- Even though we have multiple lenses, the entire optical system can be (paraxially) described using a single thin lens of some equivalent focal length and aperture number.

To make real lenses behave like ideal thin lenses, we have to use combinations of multiple lens elements (compound lenses).

Lenses - Aberrations




Lenses - Aberrations




Refraction at interfaces of complicated shapes

What shape should an interface have to make parallel rays converge to a point?




Refraction at interfaces of complicated shapes

What shape should an interface have to make parallel rays converge to a point?



Single hyperbolic interface:
point to parallel rays

Double hyperbolic interface:
point to point rays




Therefore, lenses should also have hyperbolic shapes.

Spherical lenses

In practice, lenses are often made to have spherical interfaces for ease of fabrication.

- Two roughly fitting curved surfaces ground together will eventually become spherical.




Spherical lenses don't bring parallel rays to a point.

- This is called spherical aberration.
- Approximately axial (i.e., paraxial) rays behave better.





Aberrations

Deviations from ideal thin lens behavior (e.g., imperfect focus).

- Example: spherical aberration.




Lenses - Aberrations




Oblique aberrations

These appear only as we move further from the center of the field of view.

- Contrast with spherical and chromatic, which appear everywhere.
- Many other examples (astigmatism, field curvature, etc.).



Coma







Pincushion

Barrel

Distortion example



Lenses - Aberrations




Aberrations

Deviations from ideal thin lens behavior (e.g., imperfect focus).


- Example: chromatic aberration.

focal length shifts with wavelength



glass has dispersion (refractive index changes with wavelength)

one lens cancels out dispersion of other




glasses of different refractive index


Using a doublet (two-element compound lens), we can reduce chromatic aberration.

Chromatic aberration examples






Field of View




Andrew McWilliams


Field of View




Hubble – what's the focal length?



Field of View



Hubble – what's the focal length?



A costly aberration


Hubble telescope originally suffered from severe spherical aberration.

- COSTAR mission inserted optics to correct the aberration.




Aperture

- aperture controls amount of light



Aperture

- aperture controls amount of light



Aperture size

Most lenses have variable aperture size.

- F-number notation: “f/1.4” means $f / = 1.4$ (focal length / diameter).
- Usually aperture sizes available at steps of one-half or one-third stops.
- Older lenses have separate manual aperture ring.
- Modern lenses control the aperture through a dial on the camera body (“gelled” lenses).



f/1.4



f/2.8



f/4



f/8



f/16

Aperture size

Most lenses have variable aperture size.

- F-number notation: “f/1.4” means $f / = 1.4$ (focal length / diameter).
- Usually aperture sizes available at steps of one-half or one-third stops.
- Older lenses have separate manual aperture ring.
- Modern lenses control the aperture through a dial on the camera body (“gelled” lenses).



f/1.4

f/2.8

f/4


f/8

f/16

Reminder: A “stop” changes the amount of light by a factor of 2.


Aperture

- out of focus blur




Aperture

- out of focus blur




Depth of Field

depth of field: max circle of confusion



Circle of Confusion




Circle of Confusion

$$c = M \times D \times \frac{|S - S_1|}{S}$$

Canon 5D Mark III: $f=50\text{mm}$, $f/2.8$ ($N=2.8$),

focused at 5m , pixel size= $7.5\mu\text{m}$




Hyperfocal Distance

$$H = \frac{f^2}{Nc}$$

Canon 5D Mark III: $f=50\text{mm}$, $f/2.8$ ($N=2.8$),

focused at 5m, pixel size= $7.5\mu\text{m}$




Hyperfocal Distance

$$H = \frac{f^2}{Nc}$$

Canon 5D Mark III: $f=50\text{mm}$, $f/2.8$ ($N=2.8$),

focused at 5m, pixel size= $7.5\mu\text{m}$



Depth of Field




aperture....f 1.8
shutter.....1/500
ISO.....100
distance...~3ft

aperture....f 4
shutter.....1/125
ISO.....100
distance...~3ft

aperture....f 8
shutter.....1/40
ISO.....125
distance...~3ft

Depth of Field & Motion Blur





Bokeh

artistic use



two delighted blog


coded aperture




Levin et al., SIGGRAPH 2007

Diffraction Limit


- Ernst Abbe 1873: $d = \frac{\lambda}{2n \sin \theta}$
spot radius (image space)



diffraction



Airy pattern




Diffraction Limit

- Ernst Abbe 1873:
$$d = \frac{\lambda}{2n \sin \theta} = \frac{\lambda}{2NA} \gg \frac{\lambda}{N}$$

numerical aperture

f-number
↓
N



- microscope objectives today: NA 1.4-1.6 $\rightarrow d = \lambda/2.8$
- small f-number (large NA) = high resolution but shallow depth of field
 - inherent tradeoff between “3D” information and 2D resolution
 - space-bandwidth product (uncertainty principle)

Fastest lens ever made?

Zeiss 50 mm f / 0.7 Planar lens



- Originally developed for NASA's Apollo missions.
- Stanley Kubrick somehow got to use the lens to shoot Barry Lyndon under only candlelight.

Fastest lens ever made?


Zeiss 50 mm f / 0.7 Planar lens



- Originally developed for NASA's Apollo missions.
- Stanley Kubrick somehow got to use the lens to shoot Barry Lyndon under only candlelight.

Sensors


What's a Pixel?




source: Molecular Expressions

photon to electron converter

→ photoelectric effect!



What's a Pixel?





source: Molecular Expressions

- microlens: focus light on photodiode
- color filter: select color channel
- quantum efficiency: ~50%
- fill factor: fraction of surface area used for light gathering

Two main types of imaging sensors

Two main types of imaging sensors



Charged coupled device (CCD):



- row brigade shifts charges row-by-row
- amplifiers convert charges to voltages row-by-row

Complementary metal oxide semiconductor (CMOS):

- per-pixel amplifiers convert charges to voltages
- multiplexer reads voltages row-by-row

Can you think of advantages and disadvantages of each type?

Two main types of imaging sensors



Charged coupled device (CCD):


- row brigade shifts charges row-by-row
- amplifiers convert charges to voltages row-by-row

Complementary metal oxide semiconductor (CMOS):

- per-pixel amplifiers convert charges to voltages
- multiplexer reads voltages row-by-row

Can you think of advantages and disadvantages of each type?


Two main types of imaging sensors



Charged coupled device (CCD):

- row brigade shifts charges row-by-row
- amplifiers convert charges to voltages row-by-row

- ✓ higher sensitivity
- ✓ lower noise



Complementary metal oxide semiconductor (CMOS):

- per-pixel amplifiers convert charges to voltages
- multiplexer reads voltages row-by-row

- ✓ faster read-out
- ✓ lower cost


What's a Pixel?



What's a Pixel?



What's a Pixel?





What's a Pixel?








Most Common: Color Filter Arrays

Bayer pattern





any combination possible





tradeoffs?



Assorted Pixels



- Narasimhan & Nayar @ Columbia
- multiplex anything: polarization, color, time, ND, ...



Exposure (shutter speed)

- exposure = time (e.g. 1/250, 1/60, 1, 15, bulb)



wikipedia


$\frac{1}{4}$ sec, f/3.3,



2 sec, f/6.3,

ISO (“film speed”)

sensor
sensitivity
—
analog gain applied
before ADC!





Dynamic Range

- ratio between largest and smallest possible value
- bit depth also important! common bit depths: 12-14 bits RAW / 8 bits

JPEG

high dynamic range →



Global Shutter vs. Rolling Shutter



All sensor pixels exposed at same time




Row-by-row readout of image

- shorter exposure times per pixel
- motion artifacts

What are these
dark bands?



60 Hz AC power results in 120 Hz flicker!



YouTube: user cameratest



[Sheinin et al. '17]

26 frames over 10 ms




[Sheinin et al. '17]

26 frames over 10 ms



[Sheinin et al. '17]

Photons to RAW Image




Sensor Noise

- noise is (usually) bad!
- many sources of noise: heat, electronics, amplifier gain, photon to electron conversion, pixel defects, read, ...
- different noise follows different statistical distributions, two crucial ones:
 - Gaussian
 - Poisson

Gaussian Noise


- thermal, read, amplifier
- additive, signal-independent!



+



=



Photon or Shot Noise

- signal dependent
- Poisson distribution:

$$f(k; I) = \frac{I^k e^{-I}}{k!}$$

$$s = \sqrt{I}$$

N photons: $s = \sqrt{N}$

2N photons: $s = \sqrt{2}\sqrt{N}$

nonlinear!



Signal-to-Noise Ratio (SNR)

$$SNR = \frac{\text{mean pixel value}}{\text{standard deviation of pixel value}} = \frac{m \longleftarrow \text{signal}}{S \longleftarrow \text{noise}}$$

$$= \frac{PQ_e t}{\sqrt{PQ_e t + Dt + N_r^2}}$$

P = incident photon flux (photons/pixel/sec)

Q_e = quantum efficiency

t = exposure time (sec)

D = dark current (electrons/pixel/sec), including hot pixels

N_r = read noise (rms electrons/pixel), including fixed pattern noise

Scientific Sensors

- e.g., Andor iXon Ultra 897: cooled to -100°C
- scientific CMOS & CCD
- reduce pretty much all noise, except for photon noise



Digital Photography

- optics
- aperture
- depth of field
- field of view
- exposure
- noise
- color filter arrays
- image processing pipeline




Digital Photography – Additional Resources

- What we left out: metering, autofocus, autoexposure, anti-aliasing filter, IR filter (and probably much more)
- Stanford CS 178 – Digital Photography: slides, applets, and other material online
- CMU Computational Photography 15-862
- looking for a camera? check dpreview.com

Next: The Image Processing Pipeline

- RAW images
- demosaicking
- denoising
- deblurring
- white balancing
- gamma correction
- compression



References and Further Reading

- London, Upton, Stone, “Photography”, Pearson, 11th edition, 2013
- Stanford CS 178, “Digital Photography”, Course Notes
- CMU Computational Photography course
- wikipedia