

CSC304 Lecture 15

Voting 2: Gibbard-Satterthwaite Theorem

Recap

- We introduced a plethora of voting rules
 - Plurality
 - Borda
 - Veto
 - k -Approval
 - STV
 - Plurality with runoff
 - Kemeny
 - Copeland
 - Maximin
- All these rules do something reasonable on a given preference profile
 - Only makes sense if preferences are truthfully reported

Recap

- Set of **voters** $N = \{1, \dots, n\}$
- Set of **alternatives** A , $|A| = m$
- Voter i has a **preference ranking** \succ_i over the alternatives
- **Preference profile** $\vec{\succ} =$ collection of all voter rankings
- Voting rule (social choice function) f
 - Takes as input a preference profile $\vec{\succ}$
 - Returns an alternative $a \in A$

1	2	3
a	c	b
b	a	a
c	b	c


Strategyproofness

- Would any of these rules incentivize voters to report their preferences truthfully?
- A voting rule f is **strategyproof** if for every
 - preference profiles $\vec{\succ}$,
 - voter i , and
 - preference profile $\vec{\succ}'$ such that $\succ'_j = \succ_j$ for all $j \neq i$
 - it is not the case that $f(\vec{\succ}') \succ_i f(\vec{\succ})$

Strategyproofness

- None of the rules we saw are strategyproof!
- Example: Borda Count
 - In the true profile, b wins
 - Voter 3 can make a win by pushing b to the end

	1	2	3	
	b	b	a	
Winner	a	a	b	
b	c	c	c	
	d	d	d	



	1	2	3	
	b	b	a	
	a	a	c	
Winner	c	c	d	
a	d	d	b	

Borda's Response to Critics

My scheme is
intended only for
honest men!



Random 18th
century
French dude

Strategyproofness

- Are there any strategyproof rules?
 - Sure
- Dictatorial voting rule
 - The winner is always the most preferred alternative of voter i
- Constant voting rule
 - The winner is always the same
- Not satisfactory (for most cases)



Dictatorship



Constant function

Three Properties

- **Strategyproof:** Already defined. No voter has an incentive to misreport.
- **Onto:** Every alternative can win under some preference profile.
- **Nondictatorial:** There is no voter i such that $f(\vec{\succ})$ is always the top alternative for voter i .

Gibbard-Satterthwaite

- **Theorem:** For $m \geq 3$, no deterministic social choice function can be strategyproof, onto, and nondictatorial simultaneously 😞
- **Proof:** We will prove this for $n = 2$ voters.
 - Step 1: Show that SP is equivalent to “strong monotonicity” [HW 3?]
 - **Strong Monotonicity (SM):** If $f(\vec{y}) = a$, and \vec{y}' is such that $\forall i \in N, x \in A: a \succ_i x \Rightarrow a \succ'_i x$, then $f(\vec{y}') = a$.
 - If a still defeats every alternative it defeated in every vote in \vec{y} , it should still win.

Gibbard-Satterthwaite


- **Theorem:** For $m \geq 3$, no deterministic social choice function can be strategyproof, onto, and nondictatorial simultaneously 😞
- **Proof:** We will prove this for $n = 2$ voters.
 - Step 2: Show that SP+onto implies “Pareto optimality” [HW 3?]
 - **Pareto Optimality (PO):** If $a \succ_i b$ for all $i \in N$, then $f(\vec{a}) \neq b$.
 - If there is a different alternative that *everyone* prefers, your choice is not Pareto optimal (PO).

Gibbard-Satterthwaite

- **Proof for $n=2$:** Consider problem instance $I(a, b)$

\succ_1	\succ_2
a	b
b	a
A	A
N	N
Y	Y

$I(a, b)$



\succ_1	\succ'_2
a	b
b	A
	N
	Y
A	
N	
Y	a



\succ''_1	\succ''_2
a	
	A
A	
N	
Y	
	N
	Y

Say $f(\succ_1, \succ_2) = a$

$f(\succ_1, \succ'_2) = a$

$f(\succ'') = a$

- PO: $f(\succ_1, \succ_2) \in \{a, b\}$

- PO: $f(\succ_1, \succ'_2) \in \{a, b\}$
- SP: $f(\succ_1, \succ'_2) \neq b$

- Due to strong monotonicity

Gibbard-Satterthwaite

- **Proof for $n=2$:**
 - If f outputs a on instance $I(a, b)$, voter 1 can get a elected whenever she puts a first.
 - In other words, voter 1 becomes dictatorial for a .
 - Denote this by $D(1, a)$.
 - If f outputs b on $I(a, b)$
 - Voter 2 becomes dictatorial for b , i.e., we have $D(2, b)$.
- For every $I(a, b)$, we have $D(1, a)$ or $D(2, b)$.

Gibbard-Satterthwaite

- **Proof for $n=2$:**

- On some $I(a^*, b^*)$, suppose $D(1, a^*)$ holds.
- Then, we show that voter 1 is a dictator. That is, $D(1, b)$ must hold for every other b as well.
- Take $b \neq a$. Because $|A| \geq 3$, there exists $c \in A \setminus \{a^*, b\}$.
- Consider $I(b, c)$. We either have $D(1, b)$ or $D(2, c)$.
- But $D(2, c)$ is incompatible with $D(1, a^*)$
 - Who would win if voter 1 puts a^* first and voter 2 puts c first?
- Thus, we have $D(1, b)$, as required.
- QED!

Circumventing G-S

- Randomization
 - Gibbard characterized all randomized strategyproof rules
 - Somewhat better, but still too restrictive
- Restricted preferences
 - Median for facility location (more generally, for single-peaked preferences on a line)
 - Will see other such settings later
- Money
 - E.g., VCG is nondictatorial, onto, and strategyproof, but charges payments to agents

Circumventing G-S

- Equilibrium analysis
 - Maybe good alternatives still win under Nash equilibria?
- Lack of information
 - Maybe voters don't know how other voters will vote?

Circumventing G-S

- Computational complexity (Bartholdi et al.)
 - Maybe the rule is manipulable, but it is NP-hard to find a successful manipulation?
 - Groundbreaking idea! NP-hardness can be good!!
- **Not NP-hard:** plurality, Borda, veto, Copeland, maximin, ...
- **NP-hard:** Copeland with a peculiar tie-breaking, STV, ranked pairs, ...

Circumventing G-S

- Computational complexity
 - Unfortunately, NP-hardness just says it is hard for *some worst-case instances*.
 - What if it is actually easy for most practical instances?
 - Many rules admit polynomial time manipulation algorithms for fixed #alternatives (m)
 - Many rules admit polynomial time algorithms that find a successful manipulation on almost all profiles (the fraction of profiles converges to 1)
- Interesting open problem to design voting rules that are hard to manipulate on average

Social Choice

- Let's forget incentives for now.
- Even if voters reveal their preferences truthfully, we do not have a “right” way to choose the winner.
- Who is the right winner?
 - On profiles where the prominent voting rules have different outputs, all answers seem reasonable [HW3].

Axiomatic Approach

- Define axiomatic properties we may want from a voting rule
- We already defined some:
 - Majority consistent
 - Condorcet consistent
 - Onto
 - Strategyproof
 - Strongly monotonic
 - Pareto optimal

Axiomatic Approach

- We will see four more:
 - Unanimity
 - Weak monotonicity
 - Consistency (!)
 - Independence of irrelevant alternatives (IIA)
- Problem?
 - Cannot satisfy many interesting combinations of properties
 - Arrow's impossibility result
 - Other similar impossibility results

Other Approaches?

- Statistical

- There exists an objectively true answer
 - E.g., say the question is: “Sort the candidates by the #votes they will receive in an upcoming election.”
- Every voter is trying to estimate the true ranking
- Goal is to find the most likely ground truth given votes

- Utilitarian

- Back to “numerical” welfare maximization, but we still ask voters to only report ranked preferences
 - $a \succ_i b \succ_i c$ simply means $v_i(a) \geq v_i(b) \geq v_i(c)$
- How well can we maximize welfare subject to such partial information?