

# Tutorial 5 – Kernels and Gaussian Processes

## CSC2541 Neural Net Training Dynamics – Winter 2022

Slides adapted from CSC2541: Scalable and Flexible Models of Uncertainty – Fall 2017

---

Stephan Rabanser

[stephan@cs.toronto.edu](mailto:stephan@cs.toronto.edu)



University of Toronto  
Department of Computer Science



Vector Institute for  
Artificial Intelligence


February 10, 2022

## Kernel Methods


# Recap: Basis Functions

- Basis functions allow us to use non-linear feature transformations.
- We can specify them by hand (examples below), or learn them automatically using a neural network.


$$\phi_j(x) = x^j$$



$$\phi_j(x) = \exp\left(-\frac{(x - \mu_j)^2}{2s^2}\right)$$



$$\phi_j(x) = \sigma\left(\frac{x - \mu_j}{s}\right)$$





— Bishop, Pattern Recognition and Machine Learning

# Recap: Basis Functions

- How is this useful? We can use **linear methods on non-linear features** to yield non-linear decision boundaries and regression curves.

$$\phi\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right) = \begin{bmatrix} x_1^2 \\ x_2^2 \\ \sqrt{2}x_1x_2 \end{bmatrix}$$



— <https://gregorygundersen.com/blog/2019/12/10/kernel-trick/>

## Generalized Linear Models (GLM)

- Fixed non-linear basis functions.
- Limited hypothesis space.
- Easy to optimize (convex).

## Neural Network (NN)

- Adaptive non-linear basis functions.
- Rich hypothesis space.
- Hard to optimize (non-convex).

## Towards Kernel Methods

- Feature space in GLM and NN needs to be explicitly constructed.
- Can we use a large (possibly infinite) set of fixed non-linear basis functions without explicitly constructing this space?
- Yes, by using kernel methods!

# Kernel Methods

- Kernel methods are **instance-based learners**: they assign a weight  $\theta_i$  to any training point  $\mathbf{x}_i$ .
- Predictions on new data points  $\mathbf{x}'$  make use of a **kernel function**  $\kappa(\cdot, \cdot)$  measuring the similarity of  $\mathbf{x}'$  with all points  $\mathbf{x}_i$  from the training set.
- Kernelized binary classification example:

$$\hat{y} = \text{sgn} \sum_{i=1}^n \theta_i y_i \kappa(\mathbf{x}_i, \mathbf{x}')$$

where

- $y \in \{-1, +1\}$  is the label assigned to a data point  $\mathbf{x}$ .
- $\theta_i$  is the weight for training example  $\mathbf{x}_i$ .
- $\kappa : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$  is the kernel function measuring similarity between  $\mathbf{x}, \mathbf{x}' \in \mathbb{R}$ .

# The Kernel Trick

- Let  $\phi(\cdot)$  be a set of not further specified basis functions mappings.
- Explicitly constructing a high-dimensional feature space is expensive.
- By using the **kernel trick**, we can implicitly perform operations in a high-dimensional feature space.
- In many algorithms, this **feature space only appears as a dot product**  $\langle \phi(\mathbf{x}), \phi(\mathbf{x}') \rangle = \phi(\mathbf{x})^\top \phi(\mathbf{x}')$  of input pairs  $\mathbf{x}, \mathbf{x}'$ .
- We define these dot products as the kernel function

$$\kappa(\mathbf{x}, \mathbf{x}') = \langle \phi(\mathbf{x}), \phi(\mathbf{x}') \rangle = \phi(\mathbf{x})^\top \phi(\mathbf{x}')$$

which can also be thought of as a similarity function between  $\mathbf{x}$  and  $\mathbf{x}'$ .

# Dual Representation

- Recall the regularized linear regression objective:

$$\mathcal{L}(\boldsymbol{\theta}) = \frac{1}{2} \sum_{n=1}^N (\boldsymbol{\theta}^\top \phi(\mathbf{x}_n) - y_n)^2 + \frac{\lambda}{2} \boldsymbol{\theta}^\top \boldsymbol{\theta}$$

- Finding optimal  $\boldsymbol{\theta}$ :

$$\nabla_{\boldsymbol{\theta}} \mathcal{L}(\boldsymbol{\theta}) = \sum_{n=1}^N (\boldsymbol{\theta}^\top \phi(\mathbf{x}_n) - y_n) \phi(\mathbf{x}_n) + \lambda \boldsymbol{\theta} = 0$$

$$\boldsymbol{\theta} = -\frac{1}{\lambda} \sum_{n=1}^N \underbrace{(\boldsymbol{\theta}^\top \phi(\mathbf{x}_n) - y_n)}_{\mathbf{a}_n} \phi(\mathbf{x}_n)$$

- The weights  $\boldsymbol{\theta}$  can be written as a **linear combination of the training examples**:

$$\boldsymbol{\theta} = \sum_{n=1}^N \mathbf{a}_n \phi(\mathbf{x}_n) \quad \text{where } \mathbf{a} = [\mathbf{a}_1, \dots, \mathbf{a}_n] \text{ are called the dual parameters}$$



# Dual Representation

- Substituting  $\theta$  back into linear regression  $y(\mathbf{x}) = \theta^\top \phi(\mathbf{x})$  yields:

$$\theta = \sum_{n=1}^N \mathbf{a}_n \phi(\mathbf{x}_n) \qquad y(\mathbf{x}) = \sum_{n=1}^N \mathbf{a}_n \phi(\mathbf{x}_n)^\top \phi(\mathbf{x}) = \sum_{n=1}^N \mathbf{a}_n \kappa(\mathbf{x}_n, \mathbf{x})$$


- The feature space only appears as a dot product.
- The **kernel matrix**, or **gram matrix**,  $\mathbf{K} \in \mathbb{R}^{N \times N}$  collects kernel values in a symmetric positive semi-definite matrix for all data points (**Mercer's theorem**):

$$\mathbf{K}_{ij} = \kappa(\mathbf{x}_i, \mathbf{x}_j) = \phi(\mathbf{x}_i)^\top \phi(\mathbf{x}_j)$$

- If a kernel defines such a kernel matrix, then the kernel is **valid**.


## Polynomial Kernel

$$\kappa_{\text{Pol}}(\mathbf{x}, \mathbf{x}') = (\mathbf{x}^\top \mathbf{x}' + c)^d$$



## Squared Exponential Kernel

$$\kappa_{\text{SE}}(\mathbf{x}, \mathbf{x}') = \sigma^2 \exp\left(-\frac{(\mathbf{x} - \mathbf{x}')^2}{2\ell^2}\right)$$



# Kernel Composition Rules

Let  $\kappa_1(\mathbf{x}, \mathbf{x}')$  and  $\kappa_2(\mathbf{x}, \mathbf{x}')$  be valid kernels, then the following kernels are also valid:

- $\kappa(\mathbf{x}, \mathbf{x}') = c\kappa_1(\mathbf{x}, \mathbf{x}') \quad \forall c > 0$
- $\kappa(\mathbf{x}, \mathbf{x}') = f(\mathbf{x})\kappa_1(\mathbf{x}, \mathbf{x}')f(\mathbf{x}') \quad \forall f$
- $\kappa(\mathbf{x}, \mathbf{x}') = g(\kappa_1(\mathbf{x}, \mathbf{x}')) \quad g \text{ is polynomial with coefficients } \geq 0.$
- $\kappa(\mathbf{x}, \mathbf{x}') = \exp(\kappa_1(\mathbf{x}, \mathbf{x}'))$
- $\kappa(\mathbf{x}, \mathbf{x}') = \kappa_1(\mathbf{x}, \mathbf{x}') + \kappa_2(\mathbf{x}, \mathbf{x}') \quad \text{kernel OR-ing}$
- $\kappa(\mathbf{x}, \mathbf{x}') = \kappa_1(\mathbf{x}, \mathbf{x}')\kappa_2(\mathbf{x}, \mathbf{x}') \quad \text{kernel AND-ing}$
- $\kappa(\mathbf{x}, \mathbf{x}') = \mathbf{x}^\top \mathbf{A} \mathbf{x}' \quad \mathbf{A} \text{ symmetric and p.s.d.}$

Check out the Kernel Cookbook:

<https://www.cs.toronto.edu/~duvenaud/cookbook/>


# Gaussian Processes

# Recap: Multivariate Gaussian

- Handy tool for Bayesian inference on real-valued variables
- General multivariate PDF:



$$\mathbf{x} \sim \mathcal{N}_D(\boldsymbol{\mu}, \boldsymbol{\Sigma}) = \frac{1}{\sqrt{(2\pi)^D |\boldsymbol{\Sigma}|}} e^{-\frac{1}{2}(\mathbf{x}-\boldsymbol{\mu})^\top \boldsymbol{\Sigma}^{-1}(\mathbf{x}-\boldsymbol{\mu})}$$

- Some examples of  $D = 1$  Gaussians



# Bayesian Parameter Estimation Example



Measure your heart rate at 8am



— Example from [http://videlectures.net/mlss2012\\_cunningham\\_gaussian\\_processes/](http://videlectures.net/mlss2012_cunningham_gaussian_processes/)


# Bayesian Parameter Estimation Example

Measure your heart rate at 8am and 9am



# Bayesian Parameter Estimation Example

Measuring your heart rate throughout the day





A Gaussian process describes a **distribution over functions** (infinitely long vectors).

- Notation:  $f(\mathbf{x}) \sim \mathcal{GP}(m(\mathbf{x}), \kappa(\mathbf{x}, \mathbf{x}))$
- Mean function:  $m(\mathbf{x}) = \mathbb{E}[f(\mathbf{x})]$
- Covariance function:  $\kappa(\mathbf{x}, \mathbf{x}') = \mathbb{E}[(f(\mathbf{x}) - m(\mathbf{x}))(f(\mathbf{x}') - m(\mathbf{x}'))]$

We have data points  $\mathbf{X} = [\mathbf{x}_1^\top, \dots, \mathbf{x}_n^\top]^\top$  and are interested in their function values  $f(\mathbf{X}) = (f(\mathbf{x}_1), \dots, f(\mathbf{x}_n))^\top$ .

A Gaussian process is a collection of random variables, any finite number of which have joint Gaussian distribution.

$f(\mathbf{x})$  is one such subset and has (prior) joint Gaussian distribution.

## The mean function $m$

- The mean function  $m(\cdot)$  encodes the a-priori expectation of the function.
- $m(\mathbf{x})$  will dominate the inference result in case we have not yet observed data similar to  $\mathbf{x}$ .
- Typical choice: zero-centering the data:  $m(\mathbf{x}) = 0$


## The covariance function $\kappa$

- $\kappa(\mathbf{x}, \mathbf{x}')$  measures similarity between  $\mathbf{x}$  and  $\mathbf{x}' \rightarrow$  similar data points have similar function values.
- $\kappa$  is a Mercer kernel.
- Typical choice: squared exponential kernel:  $\kappa(\mathbf{x}, \mathbf{x}') = \sigma^2 e^{-\frac{(\mathbf{x}-\mathbf{x}')^\top (\mathbf{x}-\mathbf{x}')}{2\ell^2}}$  where  $\sigma$  defines the height and  $\ell$  the width of the kernel.

# Drawing Samples From The Prior

Same procedure as for multivariate Gaussians:

1. Generate  $\mathbf{u} \in \mathbb{R}^D$  by drawing  $d$  samples from  $\mathcal{N}(\mathbf{0}, \mathbf{I}_D)$ .
2. Perform Cholesky decomposition  $\Sigma = \mathbf{L}\mathbf{L}^\top$ .
3. Compute  $\mathbf{y} = \boldsymbol{\mu} + \mathbf{L}\mathbf{u}$  where  $\mathbf{y} \sim \mathcal{N}(\boldsymbol{\mu}, \Sigma)$ .



# The Joint Distribution

We have training data  $\mathbf{X} \in \mathbb{R}^{N \times D}$ , corresponding observations  $\mathbf{y} = f(\mathbf{X})$ , and test data points  $\mathbf{X}_* \in \mathbb{R}^{N_* \times D}$  for which we want to infer function values  $\mathbf{y}_* = f(\mathbf{X}_*)$ . The GP defines the following joint distribution

$$p(\mathbf{y}, \mathbf{y}_* | \mathbf{X}, \mathbf{X}_*) = \begin{pmatrix} \mathbf{y} \\ \mathbf{y}_* \end{pmatrix} \sim \mathcal{N} \left( \begin{bmatrix} m(\mathbf{X}) \\ m(\mathbf{X}_*) \end{bmatrix}, \begin{bmatrix} \mathbf{K} + \sigma_n^2 \mathbf{I} & \mathbf{K}_* \\ \mathbf{K}_*^\top & \mathbf{K}_{**} \end{bmatrix} \right)$$

where

$$\mathbf{K} = \kappa(\mathbf{X}, \mathbf{X}) \quad \mathbf{K}_* = \kappa(\mathbf{X}, \mathbf{X}_*) \quad \mathbf{K}_{**} = \kappa(\mathbf{X}_*, \mathbf{X}_*).$$

Typically, data points are corrupted by noise  $\rightarrow$  our functions should not act as interpolators. We therefore assume

$$y_i = f(\mathbf{x}_i) + \epsilon \quad \text{where} \quad \epsilon \sim \mathcal{N}(0, \sigma_n^2).$$


# Inference with Gaussian Processes

Inferring an unknown function value and its covariance follows from conditioning multivariate Gaussians:

$$p(\mathbf{y}_* | \mathbf{y}, \mathbf{X}, \mathbf{X}_*) \sim \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$$


## Non-noisy case

- $\boldsymbol{\mu} = m(\mathbf{X}_*) + \mathbf{K}_*^\top \mathbf{K}^{-1}(\mathbf{y} - m(\mathbf{X}))$
- $\boldsymbol{\Sigma} = \mathbf{K}_{**} - \mathbf{K}_*^\top \mathbf{K}^{-1} \mathbf{K}_*$







## Noisy case

- $\boldsymbol{\mu} = m(\mathbf{X}_*) + \mathbf{K}_*^\top (\mathbf{K} + \sigma_n^2 \mathbf{I})^{-1}(\mathbf{y} - m(\mathbf{X}))$
- $\boldsymbol{\Sigma} = \mathbf{K}_{**} - \mathbf{K}_*^\top (\mathbf{K} + \sigma_n^2 \mathbf{I})^{-1} \mathbf{K}_*$




# Influence of Kernel Hyperparameters


$$\kappa(\mathbf{x}, \mathbf{x}') = \sigma^2 e^{-\frac{(\mathbf{x}-\mathbf{x}')^\top (\mathbf{x}-\mathbf{x}')}{2\ell^2}}$$



## Useful links

- <https://distill.pub/2019/visual-exploration-gaussian-processes/>
- <http://www.infinitecuriosity.org/vizgp/>
- <https://mlg.eng.cam.ac.uk/tutorials/06/es.pdf>
- [https://xavierbourretsicotte.github.io/Kernel\\_feature\\_map.html](https://xavierbourretsicotte.github.io/Kernel_feature_map.html)
- <https://www.youtube.com/watch?v=nzSBvINmg28>
- <https://www.youtube.com/watch?v=exqpaqaPG2M>

 Christopher M Bishop and Nasser M Nasrabadi, *Pattern recognition and machine learning*, vol. 4, Springer, 2006.

 Christopher K Williams and Carl Edward Rasmussen, *Gaussian processes for machine learning*, vol. 2, MIT press Cambridge, MA, 2006.